arXiv:1011.4243v1 [math.KT] 18 Nov 2010

ON KOSZULITY OF TWISTED TENSOR PRODUCTS

PASCUAL JARA, JAVIER LÓPEZ PEÑA, AND DRAGOŞ ŞTEFAN

Abstract. Let R be a semisimple ring. A pair (A, C) is called pre-Koszul if A is a connected graded R-ring and C is a compatible connected graded R-coring. To a pre-Koszul pair we associate several complexes and we show that one of them is exact if and only if all other are so. In this case we say that (A, C) is Koszul. We investigate the connection between Koszul pairs and the Koszulity of A and C. We use Koszul pairs to show that the twisted tensor product of two Koszul rings is Koszul.

Introduction

Koszul rings were introduced in the seminal paper [BGS96] as graded rings $A := \oplus_{n\geq 0} A^n$ such that: A^0 is a semisimple ring, and A^0 regarded as a graded left A-module has a resolution $P_\bullet \to A^0 \to 0$ by projective graded left A-modules, with the property that every P_n is generated by its homogeneous component of degree n. In loc. cit. the authors show that many fundamental properties of Koszul algebras hold for Koszul rings too: Koszul rings are quadratic, a ring is Koszul if and only if the Koszul complex is exact, the dual of a Koszul ring is Koszul, etc.

It is well-known that the tensor product of two Koszul algebras is Koszul too, so it is natural to ask if a similar property can be proved for Koszul rings. Of course, first of all we have to clarify what is meant by the tensor product of two graded rings. In the classical situation, when A and B are two Koszul algebras, the tensor product is performed over the base field K. Note that in this case both algebras are connected, that is $A^0 = B^0 = K$. In the general case, when A and B are just graded rings, it might happen that A^0 and B^0 to be different, so there is no good candidate for the 'base ring'. It is easy to overcome this difficulty, by fixing a semisimple ring R and dealing only with connected graded R-rings, that is graded rings with $A^0 = R$. Equivalently, a connected graded R-ring is a connected graded algebra in the monoidal category $({}_R\mathfrak{M}_R, \otimes, R)$ of R-bimodules. Clearly, if A and B are connected graded R-rings, then we may consider the graded R-bimodule $A \otimes B$ (throughout the paper the symbol $\otimes$ denotes the tensor product of two R-bimodules). Still, $A \otimes B$ do not inherit a ring structure from the existing ones of A and B, as R is not commutative. In Noncommutative Geometry a good replacement for the usual tensor product of two algebras is the twisted one, with respect to a graded twisting map $\sigma : B \otimes A \to A \otimes B$ (see §1.1 for the definition of a twisting map and of the corresponding twisted tensor product $A \otimes_\sigma B$).

In view of the foregoing, we fix a semisimple ring R, two graded R-rings A and B and a graded twisting map σ between A and B. If A and B are Koszul, now it makes sense to ask if $A \otimes_\sigma B$ is Koszul as well. The main aim of this paper is to show that this question has a positive answer, provided that the twisting map is invertible.

To approach this problem we shall use as a main tool the dual notion of R-rings, the so called R-corings. By definition, they are coalgebras in the monoidal category $({}_R\mathfrak{M}_R, \otimes, R)$. Our interest for this structure is explained by the fact that the Koszul complex of a quadratic R-ring A may be regarded in a canonical way as a connected graded R-coring C (here by connected we mean that $C^0 = R$). Moreover, the differentials of the Koszul complex may be written by using the multiplication in A and the comultiplication in C only. In the first part of the paper we show that something like the Koszul complex exists for an arbitrary pair (A, C), where A is a connected graded R-ring and C is a connected

Date: November 19, 2010.

graded R-coring that are compatible in following sense. First we ask that C^1 and A^1 are isomorphic as R-bimodules via a map $\theta_{C,A} : C^1 \to A^1$. Since both structures are graded, the multiplication in A and the comultiplication in C induce maps $m^{p,q} : A^p \otimes A^q \to A^{p+q}$ and $\Delta^{p,q} : C^{p+q} \to C^p \otimes C^q$. The second condition that we impose on A and C is that

$$m^{1,1} \circ (\theta_{C,A} \otimes \theta_{C,A}) \circ \Delta^{1,1} = 0.$$

We call such a pair pre-Koszul. In the first section of the paper we investigate the main properties of pre-Koszul pairs. We start by associating six complexes to each pre-Koszul pair (A, C). The first one is a chain complex in the category of left A-module. By symmetry we obtain another one in the category of right A-modules, and by combining these constructions we get a new chain complex in the category of A-bimodules. The other three complexes are constructed by duality. Thus, they are cochain complexes in appropriate categories of (bi)comodules.

In the case when A^1 generates A, we prove that $(A, T(A))$ is pre-Koszul, where the coring structure on $T(A) := \operatorname{Tor}^A_\bullet(R, R)$ is defined using the normalized bar resolution of $R = A^0$. Again by duality, if C is a connected graded R-coring and C is cogenerated in degree one, then $(E(C), C)$ is a pre-Koszul pair, where $E(C) := \operatorname{Ext}^\bullet_C(R, R)$.

In Theorem 2.4 we prove that one of the six complexes that are associated to a pre-Koszul pair (A, C) is exact, then the other five are also exact. In this case we say that (A, C) is a Koszul pair.

For a graded R-ring A, we prove in Theorem 2.13 that there is a graded R-coring C such that (A, C) is Koszul if and only if A is generated by A^1 and $(A, T(A))$ is Koszul, if and only if A is a Koszul ring. By duality one proves a similar result for a graded R-coring C. We end this section by showing that a component of a Koszul pair uniquely determines the other one. As a consequence of this fact,

$$E(T(A)) \cong A \qquad \text{and} \qquad T(E(C)) \cong C,$$

for any Koszul ring A and any Koszul coring C (see §2.15 for the definition of Koszul corings). Therefore $T(A)$ and $E(C)$ may be seen as the Koszul dual of A and of C, respectively. Note that these isomorphisms hold without any finiteness condition on A and C.

In the third section of the paper we prove that $A \otimes_\sigma B$ is Koszul, provided that A and B are Koszul, and that the twisting map σ is invertible. In order to do that, we consider the Koszul pairs $(A, T(A))$ and $(B, T(B))$, and show that there is a twisting map $\hat{\tau} : T(A) \otimes T(B) \to T(B) \otimes T(A)$ such that $(A \otimes_\sigma B, T(A) \otimes_{\hat{\tau}} T(B))$ is a Koszul pair.

1. Pre-Koszul pairs

In this section we introduce Koszul pairs and then we investigate their properties. First let us fix the terminology and the notation that we use. Throughout this paper R will denote a semisimple ring.

1.1. R-rings. The category of R-bimodules will be denoted by ${}_R\mathfrak{M}_R$. It is a monoidal category with respect to the tensor product of bimodules, that will be denoted by the symbol $\otimes$.

An *R-ring* is by definition an associative and unital algebra in the monoidal category ${}_R\mathfrak{M}_R$. One can see easily that an R-ring is uniquely determined by an associative and unital ring A and a morphism of unital rings $u : R \to A$. We say that an R-ring A is *graded* if it is equipped with a decomposition $A = \bigoplus_{n\in\mathbb{N}} A^n$ in ${}_R\mathfrak{M}_R$, such that the multiplication maps $A^p \otimes A^q$ to A^{p+q}. Note that the multiplication m in A induces an R-bilinear map $m^{p,q} : A^p \otimes A^q \to A$. A graded R-ring A is *connected* if $A^0 = R$. We denote by $\bar{A} = \oplus_{n>0} A^n$ the ideal generated by the homogeneous elements of positive degree. The multiplication of A induces a bimodule map $\overline{m} : \bar{A} \otimes \bar{A} \to \bar{A}$. The projection of A onto A^n will be denoted by π^n_A.

Let R^{op} be the opposite ring. If V and W are R-bimodules, then they become R^{op}-bimodules by interchanging the left and right module structures. Moreover, there is a unique isomorphism of abelian groups $\Lambda_{V,W} : V \otimes_{R^{op}} W \to W \otimes_R V$ that maps $v \otimes_{R^{op}} w$ to $w \otimes_R v$. If m denotes the multiplication

of an R-ring A, then $m^{op} := m \circ \Lambda_{A,A}$ defines a new associative multiplication on A, regarded as an R^{op}-bimodule. Together with the unit of A, the map m^{op} defines an R^{op}-ring, that will be called the *opposite R^{op}-ring* of A, and it will be denoted by A^{op}.

1.2. R-corings. An *R-coring* is a coassociative and counital coalgebra in ${}_R\mathfrak{M}_R$. Thus, an R-coring is an R-bimodule C together with a coassociative *comultiplication* $\Delta : C \to C \otimes C$ and a *counit* $\varepsilon : C \to R$, which are morphisms in ${}_R\mathfrak{M}_R$. Coassociativity can be written, using for Δ the Σ-notation $\Delta(c) := \sum c_{(1)} \otimes c_{(2)}$, as

$$\sum c_{(1)_{(1)}} \otimes c_{(1)_{(2)}} \otimes c_{(2)} = \sum c_{(1)} \otimes c_{(2)_{(1)}} \otimes c_{(2)_{(2)}},$$

for every $c \in C$. The map ε is the counit of C if, for $c \in C$,

$$\sum \varepsilon(c_{(1)})c_{(2)} = c = \sum c_{(1)}\varepsilon(c_{(2)}).$$

A *graded* R-coring is a coring (C, Δ, ε) together with a decomposition $C = \bigoplus_{n\in\mathbb{N}} C^n$ in ${}_R\mathfrak{M}_R$, such that $\Delta(C^n)$ in included into $\oplus_{p+q=n} C^p \otimes C^q$, for all n. Hence, the comultiplication of C is given by a family of R-bilinear maps $\Delta^{p,q} : C^{p+q} \to C^p \otimes C^q$. For a graded coring (C, Δ, ε) we adapt the above Σ-notation so that $\Delta^{p,q}(c)$ is written as a sum $\Delta^{p,q}(c) = \sum x^p_{(1)} \otimes x^q_{(2)}$. Therefore, in a graded coring

$$\sum \left(c^{p+q}_{(1)}\right)^p_{(1)} \otimes \left(c^{p+q}_{(1)}\right)^q_{(2)} \otimes c^r_{(2)} = \sum c^p_{(1)} \otimes \left(c^{q+r}_{(2)}\right)^q_{(1)} \otimes \left(c^{q+r}_{(2)}\right)^r_{(2)}, \tag{1}$$

for all $p, q, r \in \mathbb{N}$ and $c \in C^{p+q+r}$. It is not hard to see that, for $n > 0$, the restriction of ε to C^n is 0. In the graded case the counit satisfies the relations

$$\sum \varepsilon(c^0_{(1)})c^n_{(2)} = c = \sum c^n_{(1)}\varepsilon(c^0_{(2)}), \tag{2}$$

for every $c \in C^n$. We say that a graded R-coring C is *connected* if $C^0 = R$. We denote the projection of C onto C^n by π^n_C.

The comultiplication factors through a map $\overline{\Delta} : \overline{C} \to \overline{C} \otimes \overline{C}$, where $\overline{C} := C/C_0$ denotes the quotient R-bimodule. Note that $(\overline{C}, \overline{\Delta})$ is a coassociative coalgebra, without unit. We write p_C for the canonical projection of C onto $\overline{C}$.

If (C, Δ, ε) is an R-coring, then $\Delta^{op} := \Lambda^{-1}_{C,C} \circ \Delta$ together with ε define an R^{op}-coring C^{op}. We shall say that C^{op} is the *opposite coring* of C.

1.3. Pre-Koszul pairs. Let C be a graded R-coring and let A be a graded R-ring. The pair (A, C) is called *pre-Koszul* if A and C are connected, and there is an isomorphism $\theta_{C,A} : C^1 \to A^1$, such that

$$m^{1,1} \circ (\theta_{C,A} \otimes \theta_{C,A}) \circ \Delta^{1,1} = 0. \tag{3}$$

Using the Σ-notation, this identity is equivalent to $\sum \theta_{C,A}(c^1_{(1)})\theta_{C,A}(c^1_{(2)}) = 0$, for every $c \in C^2$.

Remark 1.4. Let (A, C) be a pre-Koszul pair, where A and C are an R-ring and an R-coring, respectively. Then (A^{op}, C^{op}) is also a pre-Koszul pair in the category of R^{op}-bimodules. Note that A^{op} and C^{op} are an R^{op}-ring and an R^{op}-coring, respectively.

1.5. The normalized bar resolution of R in $\mathfrak{M}_A$. We now want to show that, for every connected graded R-ring A, which is generated by A^1, there is a graded coring C such that (A, C) is pre-Koszul.

The groups $\mathrm{Tor}^A_\bullet(R, R)$ may be computed by using the bar resolution of R, cf. [Wei97, p. 283]. In the graded case, the normalized version of the bar resolution may be used as well.

The *right normalized bar resolution* $\bar{\beta}^r_\bullet(A)$ is the exact sequence in $\mathfrak{M}_A$

$$0 \longleftarrow R \longleftarrow A \longleftarrow \bar{A} \otimes A \longleftarrow \cdots \longleftarrow \bar{A}^{\otimes n-1} \otimes A \xleftarrow{\bar{\delta}_n} \bar{A}^{\otimes n} \otimes A \longleftarrow \cdots . \tag{4}$$

where $\bar{\delta}_0 = 0$ and

$$\bar{\delta}_n(a_1 \otimes \cdots \otimes a_n \otimes a_{n+1}) := \sum\nolimits_{i=1}^{n} (-1)^i a_1 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_{n+1}.$$

By [Wei97, p. 283], the sequence of right A-modules $\bar{\beta}^r_\bullet(A)$ is exact. Moreover, as $\bar{A}^{\otimes n}$ is projective as a right R-module, it follows that $\bar{A}^{\otimes n} \otimes A$ is projective in $\mathfrak{M}_A$. Hence $\mathrm{Tor}^A_\bullet(R,R)$ is the homology of normalized bar complex $\Omega_\bullet(A)$

$$0 \longleftarrow R \longleftarrow \bar{A} \longleftarrow \bar{A} \otimes \bar{A} \longleftarrow \cdots \longleftarrow \bar{A}^{\otimes n-1} \xleftarrow{\bar{d}_n} \bar{A}^{\otimes n} \longleftarrow \cdots, \tag{5}$$

where $\bar{d}_1 = 0$ and $\bar{d}_n$ is defined by the same formula as δ_{n-1}, for any $n > 0$ and all $a_1, \ldots, a_n \in \bar{A}$. The homology of $\Omega_\bullet(A)$ will be regarded as the definition of $\mathrm{Tor}^A_\bullet(R,R)$. The homology class of $x \in \overline{A}$ in $T^1(A)$ will be denoted by $[x]$.

The left version of the normalized bar resolution may be defined by $\bar{\beta}^l_\bullet(A) := \bar{\beta}^r_\bullet(A^{op})$. Note that $\bar{\beta}^r_n(A^{op}) \cong \overline{A}^{\otimes n} \otimes A$.

Lemma 1.6. *There is a canonical structure of differential graded R-coring on $\Omega(A) := \oplus_n \bar{A}^{\otimes n}$. In particular, $T(A) := \mathrm{Tor}^A_\bullet(R,R)$ is a connected graded R-coring, and $T^1(A)$ is the cokernel of $\overline{m}$.*

Proof. We set $\Omega_n(A) := \bar{A}^{\otimes n}$, where $\bar{A}^{\otimes 0} = R$. On $\Omega(A)$ we put the comultiplication induced by the canonical identifications $\bar{A}^{\otimes(p+q)} \simeq \bar{A}^{\otimes p} \otimes \bar{A}^{\otimes q}$. Clearly, $\Omega(A)$ becomes a connected graded R-coring. The fact that $\Omega(A)$ is a DG-coring with respect to the differentials $\bar{d}_\bullet$ is proved in [PP05, Chapter 1.1], in the particular case when R is a field and A is an R-algebra. It is easy to see that the same proof works for R-rings as well. The homology of a DG-coring has a natural structure of graded R-coring, so we conclude that $T(A)$ is a graded R-coring. This coring is connected, since $d_1 = 0$. The last assertion is obvious, as $d_2 = \overline{m}$. □

Proposition 1.7. *If A is a connected graded R-ring generated by A^1, then $(A, T(A))$ is pre-Koszul.*

Proof. We have already seen that $T(A)$ is a connected graded R-coring. Since $T^1(A) = \bar{A}/\bar{A}^2$, the canonical projection π^1_A of A onto A^1 induces a map $\theta_{T(A),A} : T^1(A) \to A^1$. It follows that $\theta_{T(A),A}$ is an isomorphism, as $\bar{A}/\bar{A}^2 = A^1$ (recall that A^1 generates A). Furthermore, an element $\omega \in T^2(A)$ can be written as the homology class of a certain $\zeta \in \operatorname{Ker} \bar{m}$. Hence $\zeta = \sum_{i=1}^n x_i \otimes y_i$, for some $x_1, \ldots, x_n$ and $y_1, \ldots, y_n$ in $\bar{A}$, such that $\sum_{i=1}^n x_i y_i = 0$. By definition,

$$\Delta^{1,1}(\omega) = \sum\nolimits_{i=1}^n [x_i] \otimes [y_i].$$

On the other hand, $x_i y_i - \pi^1(x_i)\,\pi^1(y_i)$ belongs to $\sum_{n>2} A^n$, so

$$\sum\nolimits_{i=1}^n \theta_{T(A),A}([x_i])\theta_{T(A),A}([y_i]) = \sum\nolimits_{i=1}^n \pi^1(x_i)\,\pi^1(y_i) = \pi^2\left(\sum\nolimits_{i=1}^n x_i y_i\right) = 0.$$

Thus the relation (3) holds true. □

Our goal now is to associate to a pre-Koszul pair (A, C) three cochain complexes: one in the category of graded left C-comodules and, symmetrically, one in the category of graded right C-comodules. By combining the previous two constructions, we shall get the third cochain complex, that lives in the category of graded C-bicomodules.

1.8. The categories $\mathfrak{M}^C$, ${}^C\mathfrak{M}$ and ${}^C\mathfrak{M}^C$. Let C be an R-coring. The pair (M, ρ_M) is a right C-comodule if M is a right R-module and $\rho_M : M \to M \otimes C$ is a morphisms of right R-modules such that, using the Σ-notation $\rho(m) = \sum m_{\langle 0\rangle} \otimes m_{\langle 1\rangle}$, the relations below hold true for every $m \in M$.

$$\sum m_{\langle 0\rangle_{\langle 0\rangle}} \otimes m_{\langle 0\rangle_{\langle 1\rangle}} \otimes m_{\langle 1\rangle} = \sum m_{\langle 0\rangle} \otimes m_{\langle 1\rangle_{(1)}} \otimes m_{\langle 1\rangle_{(2)}},$$

$$\sum m_{\langle 0\rangle} \varepsilon\left(m_{\langle 0\rangle}\right) = m,$$

The category $\mathfrak{M}^C$ is Grothendieck, as C is flat as a left R-module; see [Brz09, p. 264]. A right comodule M is *graded* if $M := \oplus_n M^n$, and $\rho_M(M^n) \subseteq \oplus_{p+q=n} M^p \otimes C^q$, for all n. Then the structure map of a graded comodule M is uniquely defined by a family of R-bilinear maps $\rho^{p,q}_M : M^{p+q} \to M^p \otimes C^q$.

The category ${}^C\mathfrak{M}$ of left C-comodules is constructed in a similar way. For a left C-comodule (N, ρ_N) we use the Σ notation $\rho_N(n) = \sum n_{\langle -1\rangle} \otimes n_{\langle 0\rangle}$.

A C-bicomodule is a triple (M, ρ^l_M, ρ^r_M) such that M is an R-bimodule, (M, ρ^l_M) is a left comodule, (M, ρ^r_M) is a right comodule, and for every $m \in M$

$$\sum m_{\langle -1\rangle} \otimes m_{\langle 0\rangle_{\langle 0\rangle}} \otimes m_{\langle 0\rangle_{\langle 1\rangle}} = \sum m_{\langle 0\rangle_{\langle -1\rangle}} \otimes m_{\langle 0\rangle_{\langle 0\rangle}} \otimes m_{\langle 1\rangle}.$$

A morphism of C-bicomodules is a map which is left and right C-colinear. We write ${}^C\mathfrak{M}^C$ for the category of C-bicomodules.

Lemma 1.9. *Let V be a right R-module. Then $V \otimes C$ is injective as a right C-comodule. A similar result holds for left C-comodules.*

Proof. The natural transformation

$$\Theta_{M,V} : \mathrm{Hom}_R(M, V) \to \mathrm{Hom}^C(M, V \otimes C), \quad \Theta_{M,V}(f) := (f \otimes \mathrm{Id}_C) \circ \rho \tag{6}$$

is an isomorphism for any C-comodule M. Its inverse maps a morphism of C-comodules $g : M \to V \otimes C$ to $(V \otimes \varepsilon) \circ g$. Therefore the functors $\mathrm{Hom}^C(-, V \otimes C)$ and $\mathrm{Hom}_R(-, V)$ are isomorphic. As V is injective (R is semisimple), the latter functor is exact. Then the other one is also exact. □

1.10. The complexes $\mathrm{K}^\bullet_l(A, C)$ and $\mathrm{K}^\bullet_r(A, C)$. Let (A, C) be a pre-Koszul pair. For an integer $n \geq 0$ we set

$$\mathrm{K}^n_l(A, C) := C \otimes A^n,$$

and we regard $\mathrm{K}^n_l(A, C)$ as a left C-comodule with structure map $\Delta \otimes A^n$. It is easy to see that $\mathrm{K}^n_l(A, C)$ is a graded comodule, whose homogeneous component of degree r is $C^{r-n} \otimes A^n$. The differential map $d^n_l : \mathrm{K}^n_l(A, C) \to \mathrm{K}^{n+1}_l(A, C)$ is taken to be zero on $C^0 \otimes A^n$. For $p > 0$ and $c \otimes a \in C^p \otimes A^n$,

$$d^n_l(c \otimes a) := \sum c^{p-1}_{(1)} \otimes \theta_{C,A}(c^1_{(2)})a,$$

where $\theta_{C,A} : C^1 \to A^1$, is the underlying isomorphism of the pre-Koszul pair (A, C). Note that, by definition, d^n_l maps $C^p \otimes A^n$ to $C^{p-1} \otimes A^{n+1}$, so d^n_l respects the gradings on $\mathrm{K}^n_l(A, C)$ and $K^{n+1}_l(A, C)$.

Recall that (A^{op}, C^{op}) is also a pre-Koszul pair, cf. Remark 1.4. Hence we may introduce

$$\mathrm{K}^\bullet_r(A, C) := \mathrm{K}^\bullet_l(A^{op}, C^{op}).$$

In view of the canonical isomorphism $C^{op} \otimes_{R^{op}} (A^{op})^n \cong A^n \otimes_R C$, we regard $\mathrm{K}^\bullet_r(A, C)$ as the complex whose component of degree n is the R-bimodule $\mathrm{K}^n_r(A, C) := A^n \otimes_R C$. The differential d^n_r is zero on $A^n \otimes C^0$. On the other hand, for $p > 0$ and $a \otimes c \in A^n \otimes C^p$,

$$d^n_r(a \otimes c) = \sum a\theta_{C,A}(c^1_{(1)}) \otimes c^{p-1}_{(2)}.$$

Proposition 1.11. *If (A, C) is a pre-Koszul pair then $(\mathrm{K}^\bullet_l(A, C), d^\bullet_l)$ is a complex in the category of graded left C-comodules. Analogously, $(\mathrm{K}^\bullet_r(A, C), d^\bullet_r)$ is a complex in the category of graded right C-comodules.*

Proof. We only prove the first assertion. For the second one proceeds similarly. We first claim that $d^{n+1}_l \circ d^n_l = 0$. We may assume that $p \geq 2$, otherwise the relation is trivially satisfied. Let $\theta := \theta_{C,A}$, and pick up $c \otimes a \in C^p \otimes A^n$. Since the comultiplication in C is coassociative, and using (3), we get

$$\left(d^{n+1} \circ d^n\right)(c \otimes a) = \sum (c^{p-1}_{(1)})^{p-2}_{(1)} \otimes \theta\left((c^{p-1}_{(1)})^1_{(2)}\right)\theta(c^1_{(2)})a = \sum c^{p-2}_{(1)} \otimes \theta\left((c^2_{(2)})^1_{(1)}\right)\theta\left((c^2_{(2)})^1_{(2)}\right)a = 0.$$

Hence $\mathrm{K}^\bullet_l(A, C)$ is a complex. Furthermore, for $c \otimes a \in C^p \otimes A^n$, we have

$$(C \otimes d^n)(\rho(c \otimes a)) = \sum\nolimits_{r+s=p} c^r_{(1)} \otimes d^n(c^s_{(2)} \otimes a) = \sum\nolimits_{r+s=p} \sum c^r_{(1)} \otimes (c^s_{(2)})^{s-1}_{(1)} \otimes \theta\left((c^s_{(2)})^1_{(1)}\right)a.$$

On the other hand,

$$\rho(d^n(c \otimes a)) = \rho\left(\sum c^{p-1}_{(1)} \otimes \theta(c^1_{(2)})a\right) = \sum\nolimits_{u+v=p-1} (c^{p-1}_{(1)})^u_{(1)} \otimes (c^{p-1}_{(1)})^v_{(2)} \otimes \theta(c^1_{(2)})a.$$

Taking into account that the comultiplication is coassociative, by comparing the above two relations, we deduce that d_l^n is a morphism of comodules. □

1.12. The complex $\mathrm{K}^\bullet(A,C)$. We are going to construct a cochain complex $(\mathrm{K}^\bullet(A,C),d^\bullet)$ in the category ${}^C\mathfrak{M}^C$ of C-bicomodules. By definition, $\mathrm{K}^n(A,C)=C\otimes A^n\otimes C$ and

$$d^n=d_l^n\otimes C+(-1)^{n+1}C\otimes d_r^n. \tag{7}$$

Since $d_l^\bullet$ is left C-colinear and $d_r^\bullet$ is right C-colinear it follows that $d^\bullet$ is a morphism in ${}^C\mathfrak{M}^C$. Moreover, as $d_l^{n+1}d_l^n=d_r^{n+1}d_r^n=0$, we get

$$d^{n+1}d^n=(-1)^{n+1}\left[\left(d_l^{n+1}\otimes C\right)\left(C\otimes d_r^n\right)-\left(C\otimes d_r^{n+1}\right)\left(d_l^n\otimes C\right)\right].$$

By the definition of $d_l^\bullet$ and $d_r^\bullet$, and by the fact the multiplication in A is associative, it results that $d^{n+1}d^n=0$, so $(\mathrm{K}^\bullet(A,C),d^\bullet)$ is a complex indeed.

1.13. The subcomplexes $\mathrm{K}_l^\bullet(A,C,m)$ **and** $\mathrm{K}_r^\bullet(A,C,m)$**.** For any integer $m\geq 0$ we have already noticed that d_l^n maps $C^{m-n}\otimes A^n$ to $C^{m-n-1}\otimes A^{n+1}$. Then we obtain a subcomplex $\mathrm{K}_l^\bullet(A,C,m)$ of $\mathrm{K}_l^\bullet(A,C)$ such that $\mathrm{K}_l^n(A,C,m):=C^{m-n}\otimes A^n$. Its components are trivial in degree $n>m$. Therefore, it can be displayed as follows

$$0\longrightarrow C^m\otimes A^0\longrightarrow\cdots\longrightarrow C^{m-n}\otimes A^n\xrightarrow{d_l^n}C^{m-n-1}\otimes A^{n+1}\longrightarrow\cdots\longrightarrow C^0\otimes A^m\longrightarrow 0.$$

Obviously, $\mathrm{K}_l^\bullet(A,C)=\oplus_m\mathrm{K}_l^\bullet(A,C,m)$. The complex $\mathrm{K}_r^\bullet(A,C)$ admits a similar decomposition as a direct sum of subcomplexes $\mathrm{K}_r^\bullet(A,C,m)$.

1.14. The augmented complexes $\widetilde{\mathrm{K}}_l^\bullet(A,C)$, $\widetilde{\mathrm{K}}_r^\bullet(A,C)$ **and** $\widetilde{\mathrm{K}}^\bullet(A,C)$**.** All three cochain complexes that we have introduced can be augmented in a canonical way. The inclusion of $R=C^0$ into C defines the augmentation for both $\mathrm{K}_l^\bullet(A,C)$ and $\mathrm{K}_r^\bullet(A,C)$. The augmentation map of $\mathrm{K}^\bullet(A,C)$ is, by definition, the comultiplication in C. These augmented complexes will be denoted by $\widetilde{\mathrm{K}}_l^\bullet(A,C)$, $\widetilde{\mathrm{K}}_r^\bullet(A,C)$ and $\widetilde{\mathrm{K}}^\bullet(A,C)$. Note that $\widetilde{\mathrm{K}}_l^\bullet(A,C)\cong\oplus_{m>0}\mathrm{K}_l^\bullet(A,C)\oplus\widetilde{\mathrm{K}}_l^\bullet(A,C,0)$, where $\widetilde{\mathrm{K}}_l^\bullet(A,C,0)$ is the augmented complex

$$0\longrightarrow R\xrightarrow{\cong}C^0\otimes A^m\longrightarrow 0\longrightarrow\cdots.$$

The complex $\widetilde{\mathrm{K}}_r^\bullet(A,C)$ admits a similar decomposition.

1.15. The normalized bar resolution of R **in** $\mathfrak{M}^C$**.** We are going to explain briefly how the preceding constructions and results can be dualized. Let C be an R-coring. We assume that C is connected and graded. Then R is a right C-comodule with respect to the trivial coaction. The bar resolution $\beta_r^\bullet(C)$ of R is the exact sequence

$$0\longrightarrow R\longrightarrow C\longrightarrow C\otimes C\longrightarrow\cdots\longrightarrow C^{\otimes n+1}\xrightarrow{\delta^n}C^{\otimes n+2}\longrightarrow\cdots, \tag{8}$$

where $\delta^n=\sum_{i=1}^{n+1}(-1)^{i-1}\mathrm{Id}_{C^{\otimes i-1}}\otimes\Delta\otimes\mathrm{Id}_{C^{\otimes n-i+1}}$. Note that $C^{\otimes n}\otimes C$ is an injective right C-comodule, as $C^{\otimes n}$ is injective as a right R-module. One can prove directly that this sequence is exact, by showing that $(-1)^n\mathrm{Id}_{C^{\otimes n}}\otimes\varepsilon$ is a contracting homotopy for $\beta_r^\bullet(C)$.

Recall that $\overline{C}:=C/C^0$ and that $\bar{\Delta}:\overline{C}\to\overline{C}\otimes\overline{C}$ is the unique map such that $\bar{\Delta}\circ p_C=(p_C\otimes p_C)\circ\Delta$, where p_C is the canonical projection. The *right normalized bar resolution* $\bar{\beta}_r^\bullet(C)$ of R is the resolution

$$0\longrightarrow R\longrightarrow C\longrightarrow\overline{C}\otimes C\longrightarrow\cdots\longrightarrow\overline{C}^{\otimes n}\otimes C\xrightarrow{\bar{\delta}^n}\overline{C}^{\otimes n+1}\otimes C\longrightarrow\cdots, \tag{9}$$

with $\bar{\delta}^n=\sum_{i=1}^{n}(-1)^{i-1}\mathrm{Id}_{\overline{C}^{\otimes i-1}}\otimes\bar{\Delta}\otimes\mathrm{Id}_{\overline{C}^{\otimes n-i}\otimes C}+(-1)^n\mathrm{Id}_{\bar{C}^{\otimes n}}\otimes\widetilde{\Delta}$, where $\widetilde{\Delta}:=(p_C\otimes\mathrm{Id}_C)\circ\Delta$. The left bar resolution and the left normalized bar resolution will be denoted by $\beta_l^\bullet(C)$ and $\bar{\beta}_l^\bullet(C)$, respectively.

Both resolutions $\beta_r^\bullet(C)$ and $\bar{\beta}_r^\bullet(C)$ can be used to compute $\mathrm{Ext}_C^\bullet(R,R)$. The complex that corresponds to $\bar{\beta}_r^\bullet(C)$ will play an important role in our work, so we are going to write it down explicitly. Applying $\mathrm{Hom}^C(R,-)$ to $\bar{\beta}_r^\bullet(C)$ and using the isomorphisms $\Theta_{R,-}$, we obtain the normalized bar complex $\Omega^\bullet(C)$

$$0\longrightarrow R\longrightarrow\overline{C}\longrightarrow\cdots\longrightarrow\overline{C}^{\otimes n}\xrightarrow{\bar{d}^n}\overline{C}^{\otimes n+1}\longrightarrow\cdots. \tag{10}$$

with the differential morphisms $\bar{d}_0 = 0$ and $\bar{d}^n = \sum_{i=1}^n (-1)^{i-1} \mathrm{Id}_{\overline{C}^{\otimes i-1}} \otimes \bar{\Delta} \otimes \mathrm{Id}_{\overline{C}^{\otimes n-i}}$, if $n > 0$.

Lemma 1.16. *There is a canonical connected differential graded R-ring structure on $\Omega(C) := \oplus_n \overline{C}^{\otimes n}$. In particular, $E(C) := \mathrm{Ext}_C^\bullet(R, R)$ is a connected graded R-ring, and $E^1(C) = \mathrm{Ker}\bar{\Delta}$.*

Proof. Let $\Omega^n(C) := \overline{C}^{\otimes n}$, where by convention we take $\overline{C}^{\otimes 0} = R$. Obviously, the canonical identifications $\Omega^p(C) \otimes \Omega^p(C) \cong \Omega^{p+q}(C)$ define an R-ring structure on $\Omega(C) := \oplus_n \overline{C}^{\otimes n}$ which is compatible with the differentials. □

1.17. Cogenerated corings. Let $\Delta^{p,q} : C^{p+q} \to C^p \otimes C^q$ denote the components of the comultiplication of a connected graded coring C. The maps $\Delta(n) : C^n \to C^1 \otimes \cdots \otimes C^1$ are defined inductively by

$$\Delta(n) = (\mathrm{Id}_{C^1} \otimes \Delta(n-1)) \circ \Delta^{1,n-1}, \tag{11}$$

where $\Delta(1) = \mathrm{Id}_{C^1}$ and $\Delta(2) = \Delta^{1,1}$. For $\Delta(n)$ we use the Σ-notation

$$\Delta(n)(c) = \sum c^1_{(1)} \otimes c^1_{(2)} \otimes \cdots \otimes c^1_{(n)}.$$

By the definition of $\Delta(n)$ and coassociativity, we have $\Delta(p+q) = (\Delta(p) \otimes \Delta(q)) \circ \Delta^{p,q}$, so

$$\Delta(n)(c) = \sum (c^p_{(1)})^1_{(1)} \otimes \cdots (c^p_{(1)})^1_{(p)} \otimes (c^q_{(2)})^1_{(1)} \otimes \cdots (c^q_{(2)})^1_{(q)}.$$

In the case when all $\Delta(n)$ are injective we shall say that C is *cogenerated in degree one.* Note that if C is cogenerated in degree one, then $\Delta^{p,q}$ is injective for all p and q, and conversely.

Proposition 1.18. *If C is a connected graded R-coring and C is cogenerated in degree one, then $(E(C), C)$ is a pre-Koszul pair.*

Proof. The differential of $\Omega^\bullet(C)$ is trivial in degree zero, so $E^\bullet(C)$ is connected. Let $\overline{C}^n := p_C(C^n)$. Hence $\overline{C} = \oplus_{n>0} \overline{C}^n$ and p_C is injective on each component of positive degree. In particular, $C^n \cong \overline{C}^n$.

Let $\theta : C^1 \to \overline{C}$ denote the restriction of p_C to C^1. The image of θ is included into $\mathrm{Ker}\bar{\Delta} = E^1(C)$, so we may regard θ as a map that takes values in $E^1(C)$. We claim that the pair $(E(C), C)$ satisfies the identity (3) with respect to $\theta_{C,E(C)} := \theta$. If $B^2(C)$ is the group of 2-coboundaries in the normalized bar complex $\Omega^\bullet(C)$ and $c \in C^2$, then

$$\sum p_C(c^1_{(1)}) \cdot p_C(c^1_{(2)}) = \sum p_C(c^1_{(1)}) \otimes p_C(c^1_{(2)}) + B^2 = \bar{\Delta}(p_C(c)) + B^2(C) = \bar{d}^1(p_C(c)) + B^2(C) = 0.$$

Note that the first equality is a consequence of the definition of the multiplication in $E(C)$. It remains to prove that θ is bijective. Let $\overline{c}$ be a class in $\mathrm{Ker}\bar{\Delta}$. Let c^n denote the homogeneous component of degree n of c, so $c = \sum_{n=0}^d c^n$. We claim that $c^n = 0$ for $n > 1$. We have

$$\sum_{n=2}^d \sum_{r=1}^{n-1} (p_C \otimes p_C)(\Delta^{r,n-s}(c^n)) = \bar{\Delta}(p_C(c)) = 0.$$

Since $\omega^{r,n-r} := (p_C \otimes p_C)(\Delta^{r,n-r}(c^n))$ belongs to $\overline{C}^r \otimes \overline{C}^{n-r}$, in view of the foregoing remarks, we deduce that $\omega^{r,n-r} = 0$, for every r such that $0 < r < n$. As C is flat, it follows that $p_C \otimes p_C$ is injective on $C^r \otimes C^s$. Hence $\Delta^{r,n-r}(c^n) = 0$. Since C is cogenerated in degree one we get $c^n = 0$, for any $n > 1$. In conclusion, the kernel of $\bar{\Delta}$ is included into $\overline{C}^1$. The other inclusion is obvious, so $E^1(C) = \overline{C}^1$. We deduce in particular that θ is an isomorphism. □

1.19. The cotensor product $\Box_C$. Recall that, for $X \in \mathfrak{M}^C$ and $Y \in {}^C\mathfrak{M}$, the cotensor product $X \Box_C Y$ is defined as kernel of $\rho_X \otimes \mathrm{Id}_Y - \mathrm{Id}_X \otimes \rho_V$, where ρ_X and ρ_Y are the comodule structure maps of X and Y. If C is a connected graded coring, X is a right R-module and M is a left C-comodule, then

$$(X \otimes C) \Box_C M \cong X \otimes M. \tag{12}$$

It is not difficult to show that the above isomorphism is induced by the map $\mathrm{Id}_X \otimes \varepsilon \otimes \mathrm{Id}_M$, whose inverse is $\mathrm{Id}_X \otimes \rho_M$.

In the case when C is graded and connected, we know that R is a left and right C-comodule with respect to the trivial coactions. For such a coring C, there is a canonical isomorphism $X\Box_C R \cong X^{coC}$, where the set X^{coC} of coinvariant elements contains all $x \in X$ such that $\rho_X(x) = x \otimes 1$.

Proposition 1.20. *Let (A, C) be a pre-Koszul pair. The complexes $\mathrm{K}^\bullet(A, C)\Box_C R$ and $R\Box_C \mathrm{K}^\bullet(A, C)$ are isomorphic to $\mathrm{K}^\bullet_l(A, C)$ and $\mathrm{K}^\bullet_r(A, C)$, respectively. There are similar isomorphisms for the augmented complexes.*

Proof. We have just seen that $\mathrm{K}^n(A, C)\Box_C R$ can be identified with the set of coinvariant elements in $\mathrm{K}^n(A, C)$. Clearly, a coinvariant element can be written as a sum of tensor monomials $c \otimes a \otimes 1$, with $c \in C$ and $a \in A^n$. Thus the application that maps $x \in C \otimes A^n$ to $x \otimes 1$ defines an isomorphism in ${}^C\mathfrak{M}$ between $\mathrm{K}^n(A, C)$ and $\mathrm{K}^n(A, C)\Box_C R$. For a and c as above, a straightforward computation yields

$$d^n(a \otimes c \otimes 1) = d^n_l(a \otimes c) \otimes 1.$$

Therefore, the above isomorphisms commute with the differentials of $\mathrm{K}^\bullet(A, C)\Box_C R$ and $\mathrm{K}^\bullet_l(A, C)$. □

1.21. The complexes $\mathrm{K}^l_\bullet(A, C)$, $\mathrm{K}^r_\bullet(A, C)$ and $\mathrm{K}_\bullet(A, C)$. Let (A, C) be a pre-Koszul pair. By duality, we shall construct a chain complex in each of the categories ${}_A\mathfrak{M}$, $\mathfrak{M}_A$ and ${}_A\mathfrak{M}_A$. We first construct the complex $\mathrm{K}^r_\bullet(A, C)$, whose nth component is the right A-module $\mathrm{K}^r_n(A, C) := C^n \otimes A$. The differential $d^r_n : C^n \otimes A \to C^n \otimes A$ is defined, for every $n > 0$, by the relation

$$d^r_n(c \otimes a) = \sum c^{n-1}_{(1)} \otimes \theta_{C,A}(c^1_{(2)})a.$$

Applying the previous construction to the opposite pre-Koszul pair (A^{op}, C^{op}) we obtain a complex $(\mathrm{K}^l_\bullet(A, C), d^l_\bullet)$ in ${}_A\mathfrak{M}$. Explicitly, $\mathrm{K}^l_n(A, C) := A \otimes C^n$ and

$$d^l_n(a \otimes c) = \sum a\theta_{C,A}(c^1_{(1)}) \otimes c^{n-1}_{(2)}.$$

We combine these chain complexes to obtain another one $(\mathrm{K}_\bullet(A, C), d_\bullet)$ in the category of A-bimodules. By definition, $\mathrm{K}_n(A, C) := A \otimes C^n \otimes A$. In degree n the differential of $\mathrm{K}_\bullet(A, C)$ is the map

$$d_n = d^l_n \otimes A + (-1)^n A \otimes d^r_n.$$

1.22. The subcomplexes $\mathrm{K}^r_\bullet(A, C, m)$ and $\mathrm{K}^l_\bullet(A, C, m)$. The chain complex $\mathrm{K}^r_\bullet(A, C)$ can be written as a direct sum of subcomplexes $\oplus_m \mathrm{K}^r_\bullet(A, C, m)$, where $\mathrm{K}^r_n(A, C, m) := C^{m-n} \otimes A^n$. Therefore, $\mathrm{K}^r_\bullet(A, C, m)$ is the complex

$$0 \longleftarrow C^0 \otimes A^m \longleftarrow \cdots \longleftarrow C^{n-1} \otimes A^{m-n+1} \xleftarrow{d^r_n} C^n \otimes A^{m-n} \longleftarrow \cdots \longleftarrow C^m \otimes A^0 \longleftarrow 0,$$

with the differentials inherited from $\mathrm{K}^r_\bullet(A, C)$. The complex $\mathrm{K}^l_\bullet(A, C)$ has a similar description. The corresponding subcomplexes will be denoted in this case by $\mathrm{K}^l_\bullet(A, C, m)$.

1.23. The coaugmented complexes $\widetilde{\mathrm{K}}^l_\bullet(A, C)$, $\widetilde{\mathrm{K}}^r_\bullet(A, C)$, and $\widetilde{\mathrm{K}}_\bullet(A, C)$. All chain complexes that we have introduced can be coaugmented in a canonical way. The projection of A onto $R = A^0$ defines the augmentation for $\mathrm{K}^l_\bullet(A, C)$ and $\mathrm{K}^r_\bullet(A, C)$. The augmentation map of $\mathrm{K}_\bullet(A, C)$ is given by the multiplication in A. The augmented complexes will be denoted by $\widetilde{\mathrm{K}}^l_\bullet(A, C)$, $\widetilde{\mathrm{K}}^r_\bullet(A, C)$ and $\widetilde{\mathrm{K}}_\bullet(A, C)$. Note that $\widetilde{\mathrm{K}}^l_\bullet(A, C) \cong \oplus_{m>0}\mathrm{K}^l_\bullet(A, C) \oplus \widetilde{\mathrm{K}}^l_\bullet(A, C, 0)$, where $\widetilde{\mathrm{K}}^l_\bullet(A, C, 0)$ is the coaugmented complex

$$0 \longleftarrow R \xleftarrow{\cong} C^0 \otimes A^0 \longleftarrow 0 \longleftarrow \cdots.$$

The complex $\widetilde{\mathrm{K}}^\bullet_r(A, C)$ admits a similar decomposition.

Proposition 1.24. *Let (A, C) be a pre-Koszul pair. The complexes $\mathrm{K}_\bullet(A, C) \otimes_A R$ and $R \otimes_A \mathrm{K}_\bullet(A, C)$ are isomorphic to $\mathrm{K}^l_\bullet(A, C)$ and $\mathrm{K}^r_\bullet(A, C)$, respectively. There are similar isomorphisms for the corresponding coaugmented complexes.*

Proof. Omitted, being similar to that one of Proposition 1.20. □

2. Koszul pairs

In this section we shall investigate the exactness of the augmented complex that we associated to a pre-Koszul pair (A, C). Roughly speaking, we shall show that this property is equivalent to the fact that A is a Koszul algebra and C is a Koszul coalgebra.

Proposition 2.1. *Let (A, C) be a pre-Koszul pair. The following conditions are equivalent.*

(a) The complex $\widetilde{\mathrm{K}}_l^\bullet(A, C)$ is exact.
(b) The complex $\widetilde{\mathrm{K}}_r^\bullet(A, C)$ is exact.
(c) The complex $\widetilde{\mathrm{K}}^\bullet(A, C)$ is exact.

Proof. Since $\widetilde{\mathrm{K}}_r^\bullet(A, C) \cong \widetilde{\mathrm{K}}_l^\bullet(A^{op}, C^{op})$ and $\widetilde{\mathrm{K}}^\bullet(A^{op}, C^{op}) \cong \widetilde{\mathrm{K}}^\bullet(A, C)$, it is enough to prove that (a) and (c) are equivalent. If $\widetilde{\mathrm{K}}^\bullet(A, C)$ is exact, then it is split exact in the category of right C-comodules, as its components are injective C-comodules. Thus the complex obtained by applying the functor $(-)\Box_C R$ is also exact. We conclude that (c) implies (a) by applying Proposition 1.20.

Conversely, let us assume that $\widetilde{\mathrm{K}}_l^\bullet(A, C)$ is exact. We denote, for simplicity, the complex $\widetilde{\mathrm{K}}^\bullet(A, C)$ by $\widetilde{\mathrm{K}}^\bullet$. If $X_{n,p} := C^{n-p} \otimes A^n$, then $\widetilde{\mathrm{K}}^n = \oplus_p X_{n,p} \otimes C$. Moreover, for any n and any p, the map $d_l^n \otimes \mathrm{Id}_C$ maps $X_{n,p} \otimes C$ to $X_{n+1,p} \otimes C$, while $\mathrm{Id} \otimes d_r^n$ maps the same set to $X_{n+1,p+1} \otimes C$. In particular,

$$d^n(X_{n,p} \otimes C) \subseteq X_{n+1,p} \otimes C + X_{n+1,p+1} \otimes C. \tag{13}$$

Hence $\widetilde{\mathrm{K}}_i^\bullet$ is a subcomplex of $\widetilde{\mathrm{K}}^\bullet$, where

$$\widetilde{\mathrm{K}}_i^n := \oplus_{p \geq i} X_{n,p} \otimes C.$$

We first claim that $L_i^\bullet := \widetilde{\mathrm{K}}_i^\bullet / \widetilde{\mathrm{K}}_{i+1}^\bullet$ is exact. Note that $X_{n+1,i+1} \otimes C$ is a subset of $\widetilde{\mathrm{K}}_{i+1}^{n+1}$. We have already remarked that $\mathrm{Id} \otimes d_r^n$ maps $X_{n,p} \otimes C$ to $\widetilde{\mathrm{K}}_{i+1}^{n+1}$, for every $p \geq i$. Thus the differential of $L_i^\bullet$ maps the class $x \otimes c + \widetilde{\mathrm{K}}_{i+1}^n$ in L_i^n to $d_l^n(x) \otimes c + \widetilde{\mathrm{K}}_{i+1}^{n+1}$. On the other hand, as R-bimodules, we have

$$L_i^n \cong \widetilde{\mathrm{K}}_l^\bullet(A, C, i) \otimes C.$$

In view of this isomorphism, and of the foregoing remarks, the complexes $L_i^\bullet$ and $\widetilde{\mathrm{K}}_l^\bullet(A, C, i) \otimes C$ are isomorphic. Since $\widetilde{\mathrm{K}}_l^\bullet(A, C, i)$ is a direct summand of $\widetilde{\mathrm{K}}_l^\bullet(A, C)$, it is exact. Hence $L_i^\bullet$ is exact too, as C is flat as a left R-module.

Our aim now is to show that all quotients $\widetilde{\mathrm{K}}^\bullet / \widetilde{\mathrm{K}}_i^\bullet$ are exact. For, we proceed by induction. Clearly $\widetilde{\mathrm{K}}_0^\bullet = \widetilde{\mathrm{K}}^\bullet$, so the quotient is trivially exact. Assume that $\widetilde{\mathrm{K}}^\bullet / \widetilde{\mathrm{K}}_i^\bullet$ is exact, and consider the following short exact sequence

$$0 \longrightarrow \widetilde{\mathrm{K}}_i^\bullet / \widetilde{\mathrm{K}}_{i+1}^\bullet \longrightarrow \widetilde{\mathrm{K}}^\bullet / \widetilde{\mathrm{K}}_{i+1}^\bullet \longrightarrow \widetilde{\mathrm{K}}^\bullet / \widetilde{\mathrm{K}}_i^\bullet \longrightarrow 0.$$

We have seen that $\widetilde{\mathrm{K}}_i^\bullet / \widetilde{\mathrm{K}}_{i+1}^\bullet$ is exact. Thus, a fortiori, the middle term has the same property. Now we can prove that $\widetilde{\mathrm{K}}^\bullet$ is exact. Let ω be a cocycle of degree n in $\widetilde{\mathrm{K}}^\bullet$. We choose a big enough i such that ω belongs to $M_i^n := \oplus_{p \leq i} X_{np} \otimes C$. The part of degree n of the complex $\widetilde{\mathrm{K}}^\bullet / \widetilde{\mathrm{K}}_{i+1}^\bullet$ and M_i^n are isomorphic via the canonical projection. There is a unique differential $\partial^n : M_i^n \longrightarrow M_i^{n+1}$ which commute with the projections. By construction ∂^n and d^n (the differential of $\widetilde{\mathrm{K}}^\bullet$) agree on $X_{np} \otimes C$, for any $p \leq i$. Clearly, $M_i^\bullet \cong \widetilde{\mathrm{K}}^\bullet / \widetilde{\mathrm{K}}_{i+1}^\bullet$ and ω is a cocycle in $M_i^\bullet$. Consequently, $M_i^\bullet$ is exact, and there is $\zeta \in M_i^{n-1}$ such that $\omega = \partial^n(\zeta) = d^n(\zeta)$. Thus the proposition is proved. □

Proposition 2.2. *Let $\mathrm{Hom}^C(R, \widetilde{\mathrm{K}}_r^\bullet(A, C))$ denote the complex that is obtained by applying the functor $\mathrm{Hom}^C(R, -)$ to $\widetilde{\mathrm{K}}_r^\bullet(A, C)$. Then, $\mathrm{Hom}^C(R, \widetilde{\mathrm{K}}_r^\bullet(A, C))$ is isomorphic to $(A^\bullet, 0)$.*

Proof. The map Θ_{R,A^n} is an isomorphism between $\mathrm{Hom}^C(R, \widetilde{\mathrm{K}}_r^n(A, C))$ and $\mathrm{Hom}_R(R, A^n)$. Hence we may identify $\mathrm{Hom}^C(R, \widetilde{\mathrm{K}}_r^n(A, C))$ and A^n via the map $f \mapsto (\mathrm{Id}_C \otimes \varepsilon)(f(1))$. Its inverse maps $a \in A^n$ to the C-comodule morphism f_a, which is uniquely defined by $f_a(1) = a \otimes 1$. Let ∂^n be the map that

corresponds to $\mathrm{Hom}^C(R, d_r^n)$ through the above identifications. Thus $\mathrm{Hom}^C(R, \widetilde{\mathrm{K}}_r^n(A,C)) \cong (A^\bullet, \partial^\bullet)$. For $a \in A^n$ we have

$$\partial^n(a) = (A^{n+1} \otimes \varepsilon)\left((d_r^n \circ f_a)(1)\right) = (A^{n+1} \otimes \varepsilon)(d_r^n(a \otimes 1)) = 0.$$

Therefore $\partial = 0$, so the proposition is proved. □

Proposition 2.3. *Let (A, C) be a pre-Koszul pair. The following three conditions are equivalent.*

(a) The complex $\widetilde{\mathrm{K}}_\bullet^l(A,C)$ is exact.
(b) The complex $\widetilde{\mathrm{K}}_\bullet^r(A,C)$ is exact.
(c) The complex $\widetilde{\mathrm{K}}_\bullet(A,C)$ is exact.

Proof. We have remarked that $\widetilde{\mathrm{K}}_\bullet^l(A,C)$ and $\widetilde{\mathrm{K}}_\bullet(A,C) \otimes_A R$ are isomorphic. Therefore, in view of [BG98, A3, Lemma], it results that (a) and (c) are equivalent. Analogously, (b) and (c) are equivalent. □

Theorem 2.4. *If one of the complexes $\widetilde{\mathrm{K}}_l^\bullet(A,C)$, $\widetilde{\mathrm{K}}_r^\bullet(A,C)$, $\widetilde{\mathrm{K}}^\bullet(A,C)$, $\widetilde{\mathrm{K}}_\bullet^l(A,C)$, $\widetilde{\mathrm{K}}_\bullet^r(A,C)$ or $\widetilde{\mathrm{K}}_\bullet(A,C)$ is exact, then all other five are so.*

Proof. In view of Proposition 2.1 and Proposition 2.3, it is enough to prove that $\widetilde{\mathrm{K}}_l^\bullet(A,C)$ is exact if, and only if, $\widetilde{\mathrm{K}}_\bullet^r(A,C)$ is exact. Recall that both complexes admit decompositions

$$\widetilde{\mathrm{K}}_l^\bullet(A,C) = \oplus_m \widetilde{\mathrm{K}}_l^\bullet(A,C,m) \quad \text{and} \quad \widetilde{\mathrm{K}}_\bullet^r(A,C) = \oplus_m \widetilde{\mathrm{K}}_\bullet^r(A,C,m),$$

so that for $m > 0$ the complexes $\widetilde{\mathrm{K}}_l^\bullet(A,C,m)$ and $\widetilde{\mathrm{K}}_\bullet^r(A,C,m)$ are equal to $\mathrm{K}_l^\bullet(A,C,m)$ and $\mathrm{K}_\bullet^r(A,C,m)$, respectively. By construction, they are concentrated in degree up to p, and $\mathrm{K}_l^p(A,C,m) = \mathrm{K}_{m-p}^r(A,C,m)$ and $d_l^p = d_{m-p}^r$, for any $0 \leq p \leq m$. It follows that, for such values of p,

$$\mathrm{H}^p(\mathrm{K}_l^\bullet(A,C,m)) = \mathrm{H}_{m-p}(\mathrm{K}_\bullet^r(A,C,m)).$$

All other (co)homology groups vanishes. Hence $\mathrm{K}_l^\bullet(A,C,m)$ is exact if, and only if, $\mathrm{K}_\bullet^r(A,C,m)$ is so. Clearly, $\widetilde{\mathrm{K}}_l^\bullet(A,C,0)$ and $\widetilde{\mathrm{K}}_\bullet^r(A,C,0)$ always are exact. We end the proof by remarking that a direct sum of complexes is exact if, and only if, each summand is exact. □

Corollary 2.5. *If one of the six (co)augmented complexes associated to a pre-Koszul pair (A,C) is exact then*

(i) The complex $\widetilde{\mathrm{K}}_l^\bullet(A,C)$ is a resolution of R by injective graded left C-comodules.
(ii) The complex $\widetilde{\mathrm{K}}_r^\bullet(A,C)$ is a resolution of R by injective graded right C-comodules.
(iii) The complex $\widetilde{\mathrm{K}}_\bullet^l(A,C)$ is a resolution of R by projective graded left A-modules.
(iv) The complex $\widetilde{\mathrm{K}}_\bullet^r(A,C)$ is a resolution of R by projective graded right A-modules.

Under the additional assumptions that A and C are projective R-bimodules, then

(i) If R is a separable algebra over a field K, the complex $\widetilde{\mathrm{K}}^\bullet(A,C)$ is a resolution of C by injective graded C-bicomodules.
(ii) If C is , then the complex $\widetilde{\mathrm{K}}_\bullet(A,C)$ is a resolution of A by projective graded A-bimodules.

Remark 2.6. Since R is semisimple, every R-bimodule is projective as left and right module. Though, in general, the category ${}_R\mathfrak{M}_R$ is not semisimple. For example, in the case when R is an algebra over a field K, one can see that ${}_R\mathfrak{M}_R$ is semisimple if and only if R itself is projective in this category, if and only if the multiplication map $R \otimes_K R \to R$ splits in ${}_R\mathfrak{M}_R$. By definition such an algebra is called separable. The semisimplicity of ${}^C\mathfrak{M}^C$ may be characterized in a similar way. This discussion explains why, in the above corollary, we added the assumption that A and C are projective as R-bimodules, in order that $\widetilde{\mathrm{K}}_\bullet(A,C)$ and $\widetilde{\mathrm{K}}^\bullet(A,C)$ be projective resolutions. Otherwise, we do not know if $A \otimes C^n \otimes A$ (respectively $C \otimes A^n \otimes C$) is projective (respectively injective) as an A-bimodule (respectively C-bicomodule).

Definition 2.7. A pre-Koszul pair (A,C) is called *Koszul* if the complexes in Theorem 2.4 are exact.

Corollary 2.8. *If (A,C) is Koszul then (A^{op}, C^{op}), the opposite pre-Koszul pair, is exact too.*

Proof. We have noticed that $\widetilde{\mathrm{K}}_\bullet(A^{op},C^{op})\cong\widetilde{\mathrm{K}}_\bullet(A,C)$, so one of these complexes is exact if, and only if, the other one is exact too. □

To simplify the notation, we shall write c' instead of $\pi^1_C(c)$, for all $c\in C$.

Proposition 2.9. *Let (A,C) be a pre-Koszul pair. Let $\theta:=\theta_{C,A}$. The R-bilinear maps*

$$\psi^n:\overline{C}^{\otimes n}\otimes C\to A^n\otimes C,\quad \psi^n(\overline{c}_1\otimes\cdots\otimes\overline{c}_n\otimes c)=\sum\theta(c'_1)\cdots\theta(c'_n)\otimes c$$

define a morphism $\psi^\bullet$ of chain complexes from $\overline{\beta}^\bullet_r(C)$ to $(\widetilde{\mathrm{K}}^\bullet_r(A,C),(-1)^\bullet d^\bullet_r)$.

Proof. To simplify the notation, we shall denote the homogeneous component of degree one of $c\in C$ by c'. Therefore, $c':=\pi^1_C(c)$. It is not difficult to see that the relations below hold for $x\in C$ and $c\in C^p$.

$$\sum\theta(x'_{(1)})\theta(x'_{(2)})=0\quad\text{and}\quad\sum\theta(c'_{(1)})\otimes c_{(2)}=\sum\theta(c^1_{(1)})\otimes c^{p-1}_{(2)},\tag{14}$$

Then, for $c_i\in C$, and c as before, we obtain

$$\begin{aligned}\psi^{n+1}\circ\bar{d}^n_r(\overline{c}_1\otimes\cdots\otimes\overline{c}_n\otimes c)=\sum\nolimits_{i=1}^n\sum(-1)^{i-1}\theta(c'_1)\cdots\theta((c_{i_{(1)}})')\theta((c_{i_{(2)}})')\cdots\theta(c'_n)\otimes c+\\ +(-1)^n\sum\theta(c'_1)\cdots\theta(c'_n)\theta(c'_{(1)})\otimes c_{(2)}.\end{aligned}$$

Taking into account the first relation in (14) one shows that the double sum is zero. On the other hand,

$$(-1)^nd^n_r\circ\psi^n(\overline{c}_1\otimes\cdots\otimes\overline{c}_n\otimes c)=(-1)^n\sum\theta(c'_1)\cdots\theta(c'_n)\theta(c^1_{(1)})\otimes c^{p-1}_{(2)}.$$

To conclude the proof of the proposition one uses the second identity in (14). □

Proposition 2.10. *Suppose that (A,C) is Koszul. Then A is generated in degree one and C is cogenerated in degree one. In particular, $(A,T(A))$ and $(E(C),C)$ are pre-Koszul pairs.*

Proof. In view of Theorem 2.4, we may assume that $\widetilde{\mathrm{K}}^\bullet_r(A,C)$ is exact. The part in low degrees of this sequence is

$$0\longrightarrow R\xrightarrow{\sigma^0}C\xrightarrow{d^0_r}A^1\otimes C,$$

where R is placed in degree -1. Let $\theta:=\theta_{C,A}$. By definition, d^0_r and $(\theta\otimes\mathrm{Id}_{C^n})\circ\Delta^{1,n}$ agree on C^{n+1}. As θ is bijective, $\mathrm{Ker}\Delta^{1,n}=\mathrm{Ker}d^0_r|_{C^{n+1}}$. On the other hand, since the sequence is exact in C, we get $\mathrm{Ker}d^0_r=C^0$. Then, a fortiori, $\mathrm{Ker}d^0_r|_{C^{n+1}}=0$. Thus $\Delta^{1,n}$ is injective for every n. By induction on n, using relation (11), we deduce that $\Delta(n)$ is injective, for all $n>0$.

Let us prove that A is generated by A^1. The morphism $\psi^\bullet$ that we defined in Proposition 2.9 lifts the identity of R. By hypothesis $(\widetilde{\mathrm{K}}^\bullet_r(A,C),(-1)^\bullet d^\bullet_r)$ and $\bar{\beta}^\bullet_r(C)$ are resolutions of R in $\mathfrak{M}^C$. It follows that $\lambda^\bullet:=\mathrm{Hom}^C(R,\psi^\bullet)$ is a quasi-isomorphism of complexes. On the other hand, we have proven that $\mathrm{Hom}^C(R,\widetilde{\mathrm{K}}^\bullet_r(A,C))\cong(A^\bullet,0)$. By applying the functor $\mathrm{Hom}^C(R,-)$ to $\bar{\beta}^\bullet_r(C)$, we obtain the complex $\Omega^\bullet(C)$, so $\lambda^\bullet$ may be regarded as a morphism of complexes from $\Omega^\bullet(C)$ to $(A^\bullet,0)$ satisfying

$$\lambda^n(\overline{c}_1\otimes\cdots\otimes\overline{c}_n)=\theta(c'_1)\cdots\theta(c'_n),$$

where all $\bar{c}_i$'s are in $\overline{C}$ and $c'=\pi^1(c)$, for every $c\in C$. Passing to cohomology, we get an isomorphism, $\widetilde{\lambda}^n:E^n(C)\to A^n$. Let us pick up $a\in A^n$. Then a is a cocyle in $(A^\bullet,0)$, so there is a cohomology class $\omega=\sum_{i=1}^n\overline{c}_{1,i}\otimes\cdots\otimes\overline{c}_{n,i}+B^n(C)$ in $E^n(C)$ such that $\widetilde{\lambda}^n(\omega)=a$. Equivalently,

$$a=\sum\nolimits_{i=1}^n\theta(c'_{1,i})\cdots\theta(c'_{n,i}).$$

This means that $a\in(A^1)^n$, i.e. A is generated by A^1. □

Proposition 2.11. *Let (A,C) be a pre-Koszul pair. Let $\theta:=\theta_{C,A}$. The R-bilinear maps*

$$\phi_n:A\otimes C^m\to A\otimes\overline{A}^{\otimes n},\quad \phi_n:=\mathrm{Id}_A\otimes\big(\theta^{\otimes n}\circ\Delta(n)\big)$$

define a morphism $\phi_\bullet$ of complexes from $\widetilde{\mathrm{K}}^l_\bullet(A,C)$ to the normalized bar resolution $\overline{\beta}^l_\bullet(A)$.

Proof. We first compute the value of $\phi_{n-1} \circ d_n^l$ at $a \otimes c \in A \otimes C^n$. By the definitions of ϕ_n and d_n^l, and using (11), we get

$$\phi_{n-1} \circ d_n^l(a \otimes c) = \sum a\theta(c^1_{(1)}) \otimes \theta(c^1_{(2)}) \otimes \cdots \otimes \theta(c^1_{(n)}).$$

On the other hand, using again the definitions of ϕ_n and $\overline{\delta}_n$, we get

$$\begin{aligned}\overline{\delta}^n \circ \phi_n(a \otimes c) = &\sum a\theta(c^1_{(1)}) \otimes \theta(c^1_{(2)}) \otimes \cdots \otimes \theta(c^1_{(n)}) + \\ &+ \sum_{i=1}^{n-1} \sum (-1)^i a \otimes \theta(c^1_{(1)}) \otimes \cdots \otimes \theta(c^1_{(i)})\theta(c^1_{(i+1)}) \otimes \cdots \otimes \theta(c^1_{(n)}).\end{aligned}$$

By coassociativity and the condition (3) it follows that, for a given i, the corresponding term in the above sum is zero. By comparing the results of the above computations we deduce that $\phi_\bullet$ is a morphism of cochain complexes. □

Let (A, C) be a Koszul pair. We have seen that A is generated by A^1 and $(A, T(A))$ is pre-Koszul. We are going to prove that $(A, T(A))$ is Koszul, and that this property characterizes Koszul algebras, a class of algebras that has been intensively studied in the literature; see for example [BGS96, Pri70] and the references therein.

Let V denote an R-bimodule. On $T_R(V) := \oplus_n V^{\otimes n}$ one defines a graded R-coring structure such that $\Delta^{p,q}$ is the canonical identification $V^{\otimes p+q} \cong V^{\otimes p} \otimes V^{\otimes q}$. Its counit is Id_R on $R := V^{\otimes 0}$, and zero on all other summands. To make distinction between the usual ring structure on $T_R(V)$ and the coring structure that we have just introduced, we denote them by $T^a_R(V)$ and $T^c_R(V)$, respectively.

2.12. The pre-Koszul pair (A_W, C_W). If $W \subseteq V \otimes V$ is a subbimodule, then we set $C^0_W := R$, $C^1_W := W$ and

$$C^n_W = \bigcap\nolimits_{p=0}^{n-2} V^{\otimes p} \otimes W \otimes V^{\otimes n-p-2},$$

for every $n \geq 2$. The R-bimodule $C_W := \bigoplus_{n\in\mathbb{N}} C^n_W$. We claim that C_W is a graded sub-coring of $T^c_R(V)$. To show this, we have to prove that $\Delta^{p,q}(x) \in C^p_W \otimes C^q_W$, for all p and q, and any $x \in C^{p+q}_W$. In the case when $p = 0$ or $q = 0$, this property is obvious. Let us assume that $p > 0$ and $q > 0$. Clearly $\Delta^{p,q}(x) \in C^p_R \otimes T^q$, if $p = 1$. In the case when $p \geq 2$ we have $\Delta^{p,q}(x) = x$, where in the right hand side of this identity one regards x as an element in $V^{\otimes p} \otimes V^{\otimes q}$. Thus

$$\Delta^{p,q}(x) \in \bigcap\nolimits_{i=0}^{p+q-2} V^{\otimes i} \otimes W \otimes V^{\otimes p+q-i-2} \subseteq \bigcap\nolimits_{i=0}^{p-2} V^{\otimes i} \otimes W \otimes V^{\otimes p+q-i-2} = C^p_W \otimes V^{\otimes q}.$$

Similarly, $\Delta^{p,q}(x) \in V^{\otimes p} \otimes C^q_W$. Hence $\Delta^{p,q}(x) \in C^p_W \otimes C^q_W$, that means that our claim is proved.

We define A_W to be the quotient graded R-algebra of $T^a_R(V)$ modulo the two-sided ideal generated by W. By construction, A_W and C_W are graded and connected, and $A^1_W = C^1_W$. If $x \in C^2_W = W$, then $\Delta^{1,1}(x) = x$ belongs to $V \otimes V = A^1_W \otimes A^1_W$. On the other hand the multiplication on $T^a_R(V)$ is defined by the canonical isomorphisms $V^{\otimes p} \otimes V^{\otimes q} \cong V^{\otimes p+q}$. Henceforth, $m^{1,1}(\Delta^{1,1}(x)) = x + W = 0$. Thus the conditions that are required in the definition of pre-Koszul pairs are trivially satisfied for (A_W, C_W).

Theorem 2.13. *Let A be a connected graded R-ring. The following assertions are equivalent.*

(a) The R-ring A is Koszul in the usual sense, cf. [BGS96, Definition 1.2.1].
(b) There is a graded R-coring C such that (A, C) is a Koszul pair.
(c) The R-ring A is generated by A^1 and $(A, T(A))$ is Koszul.

Proof. Let us assume that A is Koszul in the usual sense. Any Koszul algebra is quadratic, cf. [BGS96]. In fact one may choose $W \subseteq A^1 \otimes A^1$ such that $A \cong A_W$. By the very definition of the ring structure on A_W and the coring structure on C_W, it follows that $\mathrm{K}^l_\bullet(A_W, C_W)$ coincides with the Koszul complex [BGS96, p. 483]. As A is Koszul the latter complex is exact, so (A_W, C_W) is Koszul.

Let (A, C) be an arbitrary Koszul pair. We have already proven that, for such a pair, A is generated by A^1, and $(A, T(A))$ is pre-Koszul. We claim that $(A, T(A))$ is Koszul. Let $\phi_\bullet : \widetilde{\mathrm{K}}^l_\bullet(A, C) \to \overline{\beta}^l_\bullet(A)$ be the morphism of complexes that we constructed in Proposition 2.11. Proceeding as in the proof of

Proposition 2.10, one shows that $\tilde{\phi}_\bullet : (C^\bullet, 0) \to (\Omega_\bullet(A), \bar{d}_\bullet)$ is a quasi-isomorphism between the graded R-bimodule C regarded as a complex with zero differentials and $(\Omega_\bullet(A), \bar{d}_\bullet)$. This quasi-isomorphism respects the coring structures so, passing to homology, it yields an isomorphism of graded corings, still denoted $\tilde{\phi}_\bullet$, between C and $T(A)$. For $c \in C$ it verifies the relation

$$\tilde{\phi}_n(c) := \Delta(n)(c) + B_n(A),$$

where $B_n(A)$ is the group of n-boundaries in $\Omega_n(A)$. By a standard computation one checks that $\tilde{\phi}_\bullet \otimes \mathrm{Id}_C$ is an isomorphism of chain complexes from $\widetilde{\mathrm{K}}^l_\bullet(A, C)$ to $\widetilde{\mathrm{K}}^l_\bullet(A, T(A))$, so the latter complex is exact.

Finally, let as assume that $(A, T(A))$ is Koszul. Hence $\widetilde{\mathrm{K}}^l_\bullet(A, T(A))$ is a resolution of R by graded left A-modules, such that each $\widetilde{\mathrm{K}}^l_n(A, T(A))$ is generated as a module by its homogeneous component of degree n, so A is a Koszul algebra in the usual sense, cf. [BGS96, Definition 1.2.1]. □

Corollary 2.14. *If A is Koszul in the usual sense, then A^{op} is Koszul too.*

Proof. Let A be a graded ring, which is Koszul in the usual sense. Then $(A, T(A))$ is Koszul and A is generated by A^1, by the preceding theorem. Since $(A^{op}, T(A)^{op})$ is Koszul too, and A^{op} is generated by $(A^{op})^1$, it follows that A^{op} is Koszul in the usual sense. □

2.15. Koszul R-corings. The preceding theorem has a counterpart for corings. To define Koszul R-corings we dualize [BGS96, Definition 1.2.1]. Let C be an R-coring. We say that a graded right C-comodule $M := \oplus_n M^n$, with structure map ρ_M, is *cogenerated in degree n* if $\rho_M^{n,p} : M^{n+p} \to M^n \otimes C^p$ is injective for all $p \geq 0$.

The coring C is said to be (right) *Koszul* if there is an injective resolution $0 \longrightarrow R \longrightarrow I^\bullet$ of R by injective graded right C-comodules, such that each I^n is cogenerated in degree n.

Theorem 2.16. *Let C be a connected graded R-coring. The following assertions are equivalent.*

(a) The R-coring C is Koszul.
(b) There is a graded R-ring A such that (A, C) is a Koszul pair.
(c) The R-coring C is cogenerated in degree one and $(E(C), C)$ is Koszul.

Proof. By duality, from the proof of Theorem 2.13. The details are left to the reader. □

Lemma 2.17. *Let (A, C') and (A, C'') be Koszul pairs. Then $C' \cong C''$ as graded R-corings. Dually, If (A', C) and (A'', C) are Koszul, then $A' \cong A''$ as graded R-rings.*

Proof. By the proof of Theorem 2.13, there canonical coring isomorphisms $C' \cong T(A)$ and $C'' \cong T(A)$. For the proof of second part of the corollary, one uses the isomorphisms $A' \cong E(C)$ and $A'' \cong E(C)$ that are induced by $\lambda^\bullet$; see the proof of Proposition 2.10. □

Proposition 2.18. *If A is a Koszul R-coring, then $E(T(A)) \cong A$. Dually, if C is a Koszul R-coring, then $T(E(C)) \cong C$.*

Proof. Since the pair $(A, T(A))$ is Koszul, it follows that $T(A)$ is cogenerated in degree one and the pair $(E(T(A)), T(A))$ is Koszul; see the implication (b) $\Rightarrow$ (c) in Theorem 2.16. Now we can apply the previous lemma to deduce that $A \cong E(T(A))$. The second isomorphism is proved in the same way. □

Remark 2.19. Under the additional assumption that A is left finite (i.e. the homogeneous parts of A are finitely generated left R-modules), the first isomorphism in the above corollary is also proved in [BGS96, Theorem 2.10.2], in a slightly different form. More precisely, in loc. cit. the authors work with the Koszul dual of A, which is an R-ring, and not a coring.

3. The pre-Koszul pair of a twisted tensor product

Throughout R is a ring. All objects that we shall work with are R-bimodules and all maps are morphisms of R-bimodules, otherwise explicitly stated. Recall that the symbol $\otimes$ stands for the tensor product in the monoidal category of R-bimodules. In this section we consider two connected graded R-rings A and B that are generated in degree 1, and a graded twisting map (see definition below) $\sigma : B \otimes A \to A \otimes B$. Our first aim is to show that σ induces a graded twisting map of R-corings

$$\tau : T(A) \otimes T(B) \to T(B) \otimes T(A)$$

such that $(A \otimes_\sigma B, T(A) \otimes_\tau T(B))$ is a pre-Koszul pair. Of course, here $A \otimes_\sigma B$ denotes the twisted tensor product R-ring with respect to σ. Similarly, $T(A) \otimes_\tau T(B)$ is the twisted tensor product of $T(A)$ and $T(B)$ with respect to τ. Furthermore, if A and B are Koszul, then we shall show that this pair is Koszul, which in particular means that $A \otimes_\sigma B$ is a Koszul algebra. In order to prove the last mentioned result, for a twisting map σ as above, we shall also construct an invertible entwining map

$$\lambda : T(A) \otimes B \to B \otimes T(A).$$

3.1. Graded twisting maps of R-rings. Let A and B be graded R-rings. A *graded twisting map* between A and B is given by a morphisms of R-bimodules

$$\sigma : B \otimes A \to A \otimes B$$

such that $\sigma(B^p \otimes A^q) \subseteq A^q \otimes B^p$, and the following relations hold:

$$\sigma \circ (\mathrm{Id}_B \otimes m_A) = (m_A \otimes \mathrm{Id}_B) \circ (\mathrm{Id}_A \otimes \sigma) \circ (\sigma \otimes \mathrm{Id}_A), \tag{15}$$

$$\sigma \circ (m_B \otimes \mathrm{Id}_A) = (\mathrm{Id}_A \otimes m_B) \circ (\sigma \otimes \mathrm{Id}_B) \circ (\mathrm{Id}_B \otimes \sigma). \tag{16}$$

In addition, $\sigma(1_B \otimes a) = a \otimes 1_B$ and $\sigma(b \otimes 1_A) = 1_A \otimes b$, for all $a \in A$ and $b \in B$. For the restrictions of σ to $B^p \otimes A^q$ we shall use the notation $\sigma^{p,q}$.

In computations we shall denote $\sigma(b \otimes a) \in A \otimes B$, as a symbolic sum $\sum_\sigma a_\sigma \otimes b_\sigma$. Thus, for instance

$$(\mathrm{Id}_A \otimes \sigma) \circ (\sigma \otimes \mathrm{Id}_B)(b \otimes x \otimes y) = \sum\nolimits_{\sigma,\sigma'} x_\sigma \otimes y_{\sigma'} \otimes (b_\sigma)_{\sigma'} .$$

The occurrence of σ and σ' in the right hand side of the above identity indicates that the twisting map is used twice: the first time on the first and the second factors, and then once again on the second and the third factors.

If σ is a twisting map then $A \otimes B$ inherits a graded R-ring structure from the existing ones in A and B, with respect to the multiplication

$$(a' \otimes b')(a'' \otimes b'') = \sum\nolimits_\sigma a' a''_\sigma \otimes b'_\sigma b''.$$

and the unit $1_A \otimes 1_B$. The twisted tensor product will be denoted by $A \otimes_\sigma B$. In the case when R is commutative, and A and B are R-algebras, the twisted tensor product $A \otimes_\sigma B$ may be seen as a deformation of the usual tensor product algebra.

3.2. Twisting maps of DG-rings. We now assume that (A, d_A) and (B, d_B) are differential graded rings. A graded twisting map $\sigma : B \otimes A \to A \otimes B$ is called a *twisting map of DG-rings* if σ is compatible with the differential maps of A and B, in the sense that, for every q the family of morphisms $\sigma^{\bullet,q}$ is a map of complexes from $(B \otimes A^q, d_B \otimes \mathrm{Id}_{A^q})$ to $(A^q \otimes B, \mathrm{Id}_{A^q} \otimes d_B)$. Symmetrically, $\sigma^{p,\bullet}$ must be a morphism between $(B^p \otimes A, \mathrm{Id}_{B^p} \otimes d_A)$ and $(A \otimes B^p, d_A \otimes \mathrm{Id}_{B^p})$, for all p.

Proposition 3.3. *Let (A, d_A) and (B, d_B) be differential graded corings. Suppose that V is an R-bimodule.*

(i) If $\sigma_A : (V \otimes A, \mathrm{Id}_V \otimes d_A) \to (A \otimes V,\ d_A \otimes \mathrm{Id}_V)$ *is a morphism of complexes that satisfies (15), with* B *replaced by* V*, then* $\mathrm{H}_\bullet(\sigma_A)$ *still satisfies this relation (note that* $\mathrm{H}_\bullet(A)$ *is an* R*-ring, as* A *is a DG-ring). Moreover, for every* $v \in V$ *such that* $\sigma_A(v \otimes 1) = 1 \otimes v$*, then*

$$\mathrm{H}_\bullet(\sigma_A)(v \otimes \bar{1}) = \bar{1} \otimes v.$$

(ii) If $\sigma_B : (B \otimes V, d_B \otimes \mathrm{Id}_V) \to (V \otimes C, \mathrm{Id}_V \otimes d_B)$ *is a morphism of complexes that satisfies (16), with* A *replaced by* V*, then* $\mathrm{H}_\bullet(\sigma_B)$ *still satisfies this relation. Moreover, for every* $v \in V$ *such that* $\sigma_B(1 \otimes v) = v \otimes 1$*, then*

$$\mathrm{H}_\bullet(\sigma_A)(\bar{1} \otimes v) = v \otimes \bar{1}.$$

(iii) If $\sigma : B \otimes A \to A \otimes B$ *is a twisting map of differential graded corings, then* σ *induces a twisting map of corings*

$$\bar{\sigma} : \mathrm{H}^\bullet(B) \otimes \mathrm{H}^\bullet(A) \to \mathrm{H}^\bullet(B) \otimes \mathrm{H}^\bullet(A).$$

Proof. Let $(\sigma_A^p)_p$ be the components of σ. They define maps

$$\mathrm{H}^p(\sigma_A) : \mathrm{H}^p(V \otimes A) \to \mathrm{H}^p(A \otimes V).$$

By assumption every left or right R-module is flat. Hence, $\bar{\sigma} := \mathrm{H}^p(\sigma_A)$ can be seen as a map from $V \otimes \mathrm{H}^p(A)$ to $\mathrm{H}^p(A) \otimes V$. For $\omega \in \mathrm{Z}^p(A)$ let $\bar{\omega}$ denote the cohomology class of ω. If $v \in V$, then

$$\mathrm{H}^p(\sigma_A)(\bar{\omega} \otimes v) = \sum v_{\sigma_A} \otimes \overline{\omega_{\sigma_A}}, \tag{17}$$

where in the above relation we used the same notation as in the case of twisting maps of R-rings. Since σ_A satisfies the first identity in the definition of twisting maps of graded R-rings, we get

$$\sum\nolimits_{\sigma_A} \overline{(\zeta\xi)}_{\sigma_A} \otimes v_{\sigma_A} = \sum\nolimits_{\sigma,\sigma'} \overline{\zeta_{\sigma_A}\xi_{\sigma'_A}} \otimes (v_{\sigma_A})_{\sigma'_A}.$$

By the definition of the multiplication in the cohomology R-ring, and using the relation (17), we get

$$\sum\nolimits_{\bar{\sigma}} \left(\bar{\zeta}\bar{\xi}\right)_{\bar{\sigma}} \otimes v_{\bar{\sigma}} = \sum\nolimits_{\bar{\sigma},\bar{\sigma}'} \bar{\zeta}_{\bar{\sigma}}\bar{\xi}_{\bar{\sigma}'} \otimes (v_{\bar{\sigma}})_{\bar{\sigma}'}\,.$$

In conclusion, $\mathrm{H}^\bullet(\sigma_A)$ satisfies (15). Using (17), we get that $\mathrm{H}^\bullet(\sigma_A)(\bar{1} \otimes v) = v \otimes \bar{1}$, for all $v \in V$.

To prove the second part of the proposition one proceeds similarly. The third part of the proposition follows in two steps. We fix $p \geq 0$. As $\sigma^{p,\bullet}$ is a morphism of complexes from $B^p \otimes A$ to $A \otimes B^p$, that satisfies (15), by the first part of the proposition we deduce that $\mathrm{H}^\bullet(\sigma^{p,\bullet})$ has the same property. Now we consider the family of morphisms of complexes $\mathrm{H}^q(\sigma^{\bullet,q}) : B \otimes \mathrm{H}^q(A) \to \mathrm{H}^q(A) \otimes B$. One can easily see that it verifies (16), as σ does. Thus, passing to cohomology, for each p and q, we get a map

$$\bar{\sigma}^{p,q} : \mathrm{H}^p(B) \otimes \mathrm{H}^q(A) \to \mathrm{H}^q(A) \otimes \mathrm{H}^p(B).$$

The family $\bar{\sigma} = (\bar{\sigma}^{p,q})_{p,q}$, in view of the foregoing remarks, is obviously a twisting map. □

3.4. Graded twisting maps of R-corings. Let C and D be two graded R-corings. A *graded twisting map* is given by an R-bilinear map

$$\tau : C \otimes D \to D \otimes C$$

such that $\tau(C^p \otimes D^q) \subseteq D^q \otimes C^p$ and the following relations hold:

$$(\Delta_D \otimes \mathrm{Id}_C) \circ \tau = (\mathrm{Id}_D \otimes \tau) \circ (\tau \otimes \mathrm{Id}_D) \circ (\mathrm{Id}_C \otimes \Delta_D), \tag{18}$$

$$(\mathrm{Id}_D \otimes \Delta_C) \circ \tau = (\tau \otimes \mathrm{Id}_C) \circ (\mathrm{Id}_C \otimes \tau) \circ (\Delta_C \otimes \mathrm{Id}_D). \tag{19}$$

In addition, $(\mathrm{Id}_D \otimes \varepsilon_C) \circ \tau = \varepsilon_C \otimes \mathrm{Id}_D$ and $(\varepsilon_D \otimes \mathrm{Id}_C) \circ \tau = \mathrm{Id}_C \otimes \varepsilon_D$. The restrictions of τ to $C^p \otimes D^q$ will be denoted by $\tau^{p,q}$.

For a twisting map of corings we use the notation $\tau(c \otimes d) = \sum_\tau d_\tau \otimes c_\tau$, for all $c \in C$ and $d \in D$. The tensor product $C \otimes D$ inherits a graded R-coring structure, that will be denoted by $C \otimes_\tau D$. Its comultiplication Δ is defined by the formula

$$\Delta := (C \otimes \tau \otimes D) \circ (\Delta_C \otimes \Delta_D),$$

and its counit is $\varepsilon_C \otimes \varepsilon_D$. The twisted tensor product will be denoted by $C \otimes_\tau D$.

3.5. Twisting maps of *DG*-corings. We now assume that (C, d^C) and (D, d^D) are differential graded corings. A graded twisting map $\tau : C \otimes D \to D \otimes C$ is called a *twisting map of DG-corings* if τ is compatible with the differentials of C and D in the sense that $\tau^{p,\bullet}$ and $\tau^{\bullet,q}$ are morphisms of chain complexes, for all p and q.

Reasoning as in the proof of Proposition 3.3, one can show that the following result holds true.

Proposition 3.6. *Let (C, d^C) and (D, d^D) be differential graded corings. Suppose that V is an R-bimodule.*

(a) If $\tau_D : (V \otimes D, \mathrm{Id}_V \otimes d_D) \to (D \otimes V,\ d_D \otimes \mathrm{Id}_V)$ is a morphism of complexes that satisfies (18), with C replaced by V, then $\mathrm{H}_\bullet(\tau_D)$ still satisfies this relation (note that $\mathrm{H}_\bullet(D)$ is a coring, as C is a DG-coring). Moreover, if $(\varepsilon_D \otimes \mathrm{Id}_V) \circ \tau_D = \mathrm{Id}_V \otimes \varepsilon_D$, then

$$(\varepsilon_{\mathrm{H}_\bullet(D)} \otimes \mathrm{Id}_V) \circ \mathrm{H}_\bullet(\tau_D) = \mathrm{Id}_V \otimes \varepsilon_{\mathrm{H}_\bullet(D)}.$$

(b) If $\tau_C : (C \otimes V, d_C \otimes \mathrm{Id}_V) \to (V \otimes C, \mathrm{Id}_V \otimes d_C)$ is a morphism of complexes that satisfies (19), with D replaced by V, then $\mathrm{H}_\bullet(\tau_C)$ still satisfies this relation. Moreover, if $(\mathrm{Id}_V \otimes \varepsilon_C) \circ \tau_C = \varepsilon_C \otimes \mathrm{Id}_V$, then

$$(\mathrm{Id}_V \otimes \varepsilon_{\mathrm{H}_\bullet(C)}) \circ \mathrm{H}_\bullet(\tau_C) = \mathrm{Id}_V \otimes \varepsilon_{\mathrm{H}_\bullet(C)}.$$

(c) If $\tau : C \otimes D \to D \otimes C$ is a twisting map of differential graded corings, then τ induces a twisting map of corings

$$\bar{\tau} : \mathrm{H}_\bullet(C) \otimes \mathrm{H}_\bullet(D) \to \mathrm{H}_\bullet(D) \otimes \mathrm{H}_\bullet(C).$$

3.7. Graded entwining maps. Let A be a graded R-ring, and let C be a graded R-coring. Let

$$\lambda : C \otimes A \to A \otimes C$$

be an R-bilinear map such that $\lambda(C^p \otimes A^q) \subseteq A^q \otimes C^p$. We denote the restriction of λ to $C^p \otimes A^q$ by $\lambda^{p,q}$. We say that λ is a *graded entwining map* if $\lambda(c \otimes 1_A) = 1_A \otimes c$ and $(\mathrm{Id}_A \otimes \varepsilon_C) \circ \lambda = \varepsilon_C \otimes \mathrm{Id}_A$, and the following two relations hold:

$$\lambda \circ (C \otimes m_A) = (m_A \otimes C) \circ (A \otimes \lambda) \circ (\lambda \otimes A), \tag{20}$$

$$(A \otimes \Delta_C) \circ \lambda = (\lambda \otimes C) \circ (C \otimes \lambda) \circ (\Delta_C \otimes A). \tag{21}$$

Let $\lambda : C \otimes A \to A \otimes C$ be a graded entwining map. We assume that, in addition, A is a differential graded R-ring and C is a differential graded R-coring. We say that a graded entwining map λ is a differential graded entwining map if $\lambda^{p,\bullet}$ and $\lambda^{\bullet,q}$ are morphisms of chain complexes, for any p and q. We state for future reference, without proof, the following proposition.

Proposition 3.8. *If (A, d^A) is a DG-ring, (C, d^C) is a DG-coring and $\lambda : C \otimes A \to A \otimes C$ is a differential graded entwining map, then λ induces an graded entwining map*

$$\bar{\lambda} : \mathrm{H}_\bullet(C) \otimes \mathrm{H}^\bullet(A) \to \mathrm{H}^\bullet(A) \otimes \mathrm{H}_\bullet(C).$$

Recall that, for every R-bimodule V, the graded R-ring $T^a_R(V)$ and the graded R-coring $T^c_R(V)$ have the same homogeneous component of degree n, namely $V^{\otimes n}$. The multiplication and the comultiplication are induced by the canonical identification $V^{\otimes p} \otimes V^{\otimes p} \cong V^{\otimes p+q}$.

Let $\alpha : W \otimes V \to V \otimes W$ be a morphism of R-bimodules. We define $\alpha^{p,q} : W^{\otimes p} \otimes V^{\otimes q} \to V^{\otimes q} \otimes W^{\otimes p}$ as follows. First, we take $\alpha^{0,p}$ to be the canonical identification $R \otimes V^{\otimes p} \cong V^{\otimes p} \otimes R$, and then we define $\alpha^{q,0}$ analogously. For $q > 0$ we set

$$\alpha^{1,q} := (V^{\otimes q-1} \otimes \alpha) \circ (V^{\otimes q-2} \otimes \alpha \otimes V) \circ \cdots \circ (\alpha \otimes V^{\otimes q-1}).$$

Note that $\alpha^{1,1} = \alpha$. Finally, if $p > 0$ we put

$$\alpha^{p,q} := \left(\alpha^{1,q} \otimes W^{\otimes p-1}\right)\left(W \otimes \alpha^{1,q} \otimes W^{\otimes p-2}\right) \circ \cdots \circ \left(W^{\otimes p-1} \otimes \alpha^{1,q}\right).$$

Let $\tilde{\alpha} : T_R(W) \otimes T_R(V) \to T_R(V) \otimes T_R(W)$ denote the unique map induced by the family $(\alpha^{p,q})_{p,q\in\mathbb{N}}$. Here $T_R(V)$ and $T_R(W)$ are regarded just as R-bimodules, without any other additional structure.

The map $\tilde{\alpha}$ provides important examples of twisting and entwining maps. The next result is 'folklore', so we omit the proof.

Lemma 3.9. *Let $\alpha : W \otimes V \to V \otimes W$ be a morphism of R-bimodule. Let $\tilde{\alpha}$ be the lifting of α as above.*

(a) The map $\tilde{\alpha} : T^a_R(W) \otimes T^a_R(V) \to T^a_R(V) \otimes T^a_R(W)$ is a graded twisting map.
(b) The map $\tilde{\alpha} : T^c_R(W) \otimes T^c_R(V) \to T^c_R(V) \otimes T^c_R(W)$ is a graded twisting map.
(c) The map $\tilde{\alpha} : T^c_R(W) \otimes T^a_R(V) \to T^a_R(V) \otimes T^c_R(W)$ is a graded entwining map.

Let $\sigma : B \otimes A \to A \otimes B$ be a twisting map of graded connected R-rings. Recall that $\Omega^\bullet(A)$ is a DG-coring, and $\Omega(A) \cong T^c_R(\bar{A})$ as graded corings. Since σ respects the gradings on A and on B, it maps $\overline{B} \otimes \bar{A}$ to $\bar{A} \otimes \overline{B}$. Let α denote the restriction of σ to $\overline{B} \otimes \bar{A}$. If $(\alpha^{p,q})_{p,q}$ are the liftings of α as above, then $\tilde{\alpha} : \Omega(B) \otimes \Omega(A) \to \Omega(A) \otimes \Omega(B)$ is a graded twisting map of R-corings. Using the relations

$$\alpha^{p+q,r} = (\overline{B}^{\otimes p+1} \otimes \alpha^{q-1,r}) \circ (\overline{B}^{\otimes p-1} \otimes \alpha^{2,r} \otimes \overline{B}^{\otimes q-1}) \circ (\alpha^{p-1,r} \otimes \overline{B}^{q+1}),$$

together with the fact that σ is a twisting map of R-rings, one shows by induction that $\alpha^{\bullet,q}$ is a morphism of complexes. In a similar way it follows that $\alpha^{\bullet,q}$ is morphism of complexes too. Therefore $\tilde{\alpha}$ is a twisting map of DG-corings, so we may apply the Proposition 3.6 for $\tilde{\alpha}$ to get a twisting map of corings $\tau : T(B) \otimes T(A) \to T(A) \otimes T(B)$.

For $q > 0$, let $\beta^{p,q}$ be the restriction of $\alpha^{p,1}$ to $\overline{B}^{\otimes p} \otimes A^q$. We take $\beta^{p,0}$ to be the canonical identification $\overline{B}^{\otimes p} \otimes R \cong R \otimes \overline{B}^{\otimes p}$. Since $(\beta^{p,q})_{p,q}$ defines an entwining map between $\Omega(B)$ and A, which is compatible with the differential maps (one regards A as a chain complex with trivial differentials), by Proposition 3.8 there is a graded entwining map $\lambda : T(B) \otimes A \to A \otimes T(B)$. Note that, by construction

$$\left(\theta_{T(A),A} \otimes \mathrm{Id}_{T^p(B)}\right) \circ \tau^{p,1} = \lambda^{p,1} \circ (\mathrm{Id}_{T^p(B)} \otimes \theta_{T(A),A}), \tag{22}$$

$$(\mathrm{Id}_{A^q} \otimes \theta_{T(B),B}) \circ \lambda^{1,q} = \sigma^{1,q} \circ (\theta_{T(B),B} \otimes \mathrm{Id}_{A^q}), \tag{23}$$

where $\theta_{T(A),}$ and $\theta_{T(B),B}$ are the isomorphisms corresponding to the pre-Koszul pairs $(A, T(A))$ and $(B, T(B))$, respectively. In particular, taking $p = q = 1$ and composing the second relation to the right with $\mathrm{Id}_{T^1(B)} \otimes \theta_A$ we deduce that

$$\sigma^{1,1} \circ (\theta_{T(B),B} \otimes \theta_{T(A),A}) = (\theta_{T(A),A} \otimes \theta_{T(B),B}) \circ \tau^{1,1}. \tag{24}$$

Of course, if σ is an invertible twisting map, then τ and λ are invertible too. Since σ^{-1} is a twisting map we can use it to produce a new invertible graded twisting map of corings $\tau' : T(A)\otimes T(B) \to T(B)\otimes T(A)$ and new graded invertible entwining map $\lambda' : T(A) \otimes B \to B \otimes T(A)$. Clearly, τ' and λ' satisfy the above relations with A and B interchanged, and with $\sigma^{1,1}$ and $\sigma^{1,q}$ replaced by $\left(\sigma^{1,1}\right)^{-1}$ and $\left(\sigma^{q,1}\right)^{-1}$, respectively. Later on in the paper, we shall use only σ, τ' and λ'. For this reason we shall change the notation, writing τ and λ instead of τ' and λ', respectively. We summarize up the foregoing remarks in the next proposition.

Proposition 3.10. *Let $\sigma : B \otimes A \to A \otimes B$ be an invertible twisting map of connected graded R-corings. If A and B are generated in degree one, then there are an invertible graded twisting map $\tau : T(A)\otimes T(B) \to T(B)\otimes T(A)$ and an invertible graded entwining map $\lambda : T(A)\otimes B \to B\otimes T(A)$ such that the relations (22), (23) and (24) hold with A and B interchanged, and with $\sigma^{1,1}$ and $\sigma^{1,q}$ replaced by $\left(\sigma^{1,1}\right)^{-1}$ and $\left(\sigma^{q,1}\right)^{-1}$, respectively.*

Our goal is to construct a pre-Koszul pair $(A \otimes_\sigma B, T(A) \otimes_{\hat{\tau}} T(B))$, where $\hat{\tau}$ and τ are equal up to a sign (see Remark 3.12 for the definition of $\hat{\tau}$). Then we shall prove that this pair is Koszul, provided that the algebras A and B are Koszul algebras. In fact we shall do that in a more general setting, that we are now going to introduce.

3.11. Notation and assumptions. From now on, (A, C) and (B, D) are two pre-Koszul pairs that are equipped with the following extra data:

(i) An invertible graded twisting map $\sigma : B \otimes A \to A \otimes B$.
(ii) An invertible graded twisting map $\tau : C \otimes D \to D \otimes C$.
(iii) An invertible entwining map $\lambda : C \otimes B \to B \otimes C$

We impose on these data the following two conditions:

$$\lambda^{p,1} \circ (\mathrm{Id}_{C^p} \otimes \theta_{D,B}) = (\theta_{D,B} \otimes \mathrm{Id}_{C^p}) \circ \tau^{p,1}, \tag{25}$$

$$\left(\sigma^{q,1}\right)^{-1} \circ (\theta_{C,A} \otimes \mathrm{Id}_{B^q}) = (\mathrm{Id}_{B^q} \otimes \theta_{C,A}) \circ \lambda^{1,q}, \tag{26}$$

where $\theta_{C,A} : C^1 \to A^1$ and $\theta_{D,B} : D^1 \to A^1$ are the isomorphisms corresponding to the pre-Koszul pairs (A, C) and (B, D), respectively.

Note that, for an invertible twisting map $\sigma : B \otimes A \to A \otimes B$, the pairs $(A, T(A))$ and $(B, T(B))$ fulfil the conditions (25)–(26), where τ and λ are as in Proposition 3.10.

Let us remark that the identities (25) and (26) together yields

$$\sigma^{1,1} \circ (\theta_{D,B} \otimes \theta_{C,A}) \circ \tau^{1,1} = \theta_{C,A} \otimes \theta_{D,B}.$$

Equivalently, for $c \in C^1$ and $d \in D^1$, we have

$$\sum\nolimits_{\sigma,\tau} \theta_{C,A}(c_\tau)_\sigma \otimes \theta_{D,B}(d_\tau)_\sigma = \theta_C(c) \otimes \theta_D(d). \tag{27}$$

Remark 3.12. If $\tau : C \otimes D \to D \otimes C$ is a twisting map of graded coalgebras, then the map $\hat{\tau} : C \otimes D \to D \otimes C$ defined by $\hat{\tau}^{p,q} := (-1)^{pq}\tau^{p,q}$ is also a graded twisting map of coalgebras, to which we can associate the twisted tensor product $C \otimes_{\hat{\tau}} D$.

Proposition 3.13. *Keeping the notation and assumptions in* §3.11*, then the pair* $(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$ *is pre-Koszul.*

Proof. It is obvious that $A \otimes_\sigma B$ and $C \otimes_{\hat{\tau}} D$ are connected. Also, by definition we have

$$(A \otimes_\sigma B)^1 = (A^0 \otimes B^1) \oplus (A^1 \otimes B^0) = (R \otimes B^1) \oplus (A^1 \otimes R),$$
$$(C \otimes_{\hat{\tau}} D)^1 = (C^0 \otimes D^1) \oplus (C^1 \otimes D^0) = (R \otimes D^1) \oplus (C^1 \otimes R).$$

Then we define $\theta : (C \otimes_{\hat{\tau}} D)^1 \to (A \otimes_\sigma B)^1$ by $\theta = (\theta_{C,A} \otimes \mathrm{Id}_R) \oplus (\mathrm{Id}_R \otimes \theta_{D,B})$. Let $\cdot_\sigma$ denote the multiplication on $A \otimes_\sigma B$. It remains to show that, for any $x \in (C \otimes_{\hat{\tau}} D)^2$,

$$\sum \theta(x^1_{(1)}) \cdot_\sigma \theta(x^1_{(2)}) = 0. \tag{28}$$

Let $\Delta := \Delta_{C \otimes_{\hat{\tau}} D}$. First, recall that $\Delta^{1,1} : (C \otimes_{\hat{\tau}} D)^2 \to (C \otimes_{\hat{\tau}} D)^1 \otimes (C \otimes_{\hat{\tau}} D)^1$, where

$$(C \otimes_{\hat{\tau}} D)^2 = (C^2 \otimes R) \oplus (C^1 \otimes D^1) \oplus (R \otimes D^2).$$

Therefore, we can restrict ourselves to prove (28) for x belonging to each of the former three direct summands. Let us consider in the first place the case $x \in C^2 \otimes R$, so we can write $x = c \otimes 1$ for $c \in C^2$. By definition of the comultiplication on $C \otimes_{\hat{\tau}} D$, we have

$$\Delta^{1,1}\big|_{C^2 \otimes R} = (C \otimes \hat{\tau}^{1,0} \otimes R) \circ (\Delta_C^{1,1} \otimes \Delta_D^{0,0}),$$

and since $\hat{\tau}|_{C \otimes R}$ is just the flip, we get

$$\Delta^{1,1}(c \otimes 1) = \sum (c^1_{(1)} \otimes 1) \otimes (c^1_{(2)} \otimes 1).$$

Henceforth, for $x = c \otimes 1 \in C^2 \otimes R$ we have

$$\sum \theta(x^1_{(1)}) \cdot_\sigma \theta(x^1_{(2)}) = \sum (\theta_{C,A}(c^1_{(1)}) \otimes 1) \cdot_\sigma (\theta_{C,A}(c^1_{(2)}) \otimes 1) = \sum \theta_{C,A}(c^1_{(1)})\theta_{C,A}(c^1_{(2)}) \otimes 1 = 0,$$

since (A, C) is a pre-Koszul pair. If $x \in C^0 \otimes D^2$, the computations are done in a similar way.

Let us finally assume that $x \in C^1 \otimes D^1$. We can just prove the condition for a generator of the form $x = c \otimes d$ with $c \in C^1$, $d \in D^1$. Recall that, for any graded coring C and $c \in C^1$ we have $\Delta c = 1 \otimes c + c \otimes 1$. Thus

$$\begin{aligned}\Delta(c \otimes d) &= (C_1 \otimes \hat{\tau} \otimes C_2)(\Delta_C(c) \otimes \Delta_D(d)) \\ &= (c \otimes d) \otimes (1 \otimes 1) + (c \otimes 1) \otimes (1 \otimes d) + \ (1 \otimes 1) \otimes (c \otimes d) + 1 \otimes \hat{\tau}^{1,1}(c \otimes d) \otimes 1 \\ &= (c \otimes d) \otimes (1 \otimes 1) + (c \otimes 1) \otimes (1 \otimes d) + \ (1 \otimes 1) \otimes (c \otimes d) - 1 \otimes \tau^{1,1}(c \otimes d) \otimes 1,\end{aligned}$$

where in the last step we used that $\hat{\tau}(c \otimes d) = -\tau(c \otimes d)$. The component of the latest expression belonging to $(C \otimes_{\hat{\tau}} D)^1 \otimes (C \otimes_{\hat{\tau}} D)^1$ is precisely

$$\Delta^{1,1}(c \otimes d) = (c \otimes 1) \otimes (1 \otimes d) - 1 \otimes \tau^{1,1}(c \otimes d) \otimes 1 = (c \otimes 1) \otimes (1 \otimes d) - \sum\nolimits_{\tau} (1 \otimes d_\tau) \otimes (c_\tau \otimes 1).$$

Henceforth, composing with the product we get

$$\begin{aligned}\sum \theta(x^1_{(1)}) \cdot_\sigma \theta(x^1_{(2)}) &= \theta(c \otimes 1) \cdot_\sigma \theta(1 \otimes d) - \sum\nolimits_{\tau} \theta(1 \otimes d_\tau) \cdot_\sigma \theta(c_\tau \otimes 1) \\ &= \theta_{C,A}(c) \otimes \theta_{D,B}(d) - \sum\nolimits_{\tau} 1 \otimes \theta_{D,B}(d_\tau) \cdot_\sigma \theta_{C,A}(c_\tau) \otimes 1 \\ &= \theta_{C,A}(c) \otimes \theta_{D,B}(d) - \sum\nolimits_{\sigma,\tau} \theta_{C,A}(c_\tau)_\sigma \otimes \theta_{D,B}(d_\tau)_\sigma \\ &= \theta_{C,A}(c) \otimes \theta_{D,B}(d) - \theta_{C,A}(c) \otimes \theta_{D,B}(d) = 0.\end{aligned}$$

where for the penultimate equality we used (27). □

Theorem 3.14. *Keeping the notation and the assumptions in §3.11, then $K^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$ is the tensor product of the complexes $K^l_\bullet(A, C)$ and $K^l_\bullet(B, D)$. In the particular case when (A, C) and (B, D) are Koszul, it follows that $(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$ is Koszul two.*

Proof. In Proposition 3.13 we have already proven that $(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$ is a pre-Koszul pair. Then it makes sense to consider the complex $K_\bullet := K^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$. Let us show that $K_\bullet$ is isomorphic to the tensor product of $K'_\bullet := K^l_\bullet(A, C)$ and $K''_\bullet := K^l_\bullet(B, D)$. Let

$$\partial_n : (A \otimes_\sigma B) \otimes (C \otimes_{\hat{\tau}} D)^n \to (A \otimes_\sigma B) \otimes (C \otimes_{\hat{\tau}} D)^{n-1}$$

denote the differential map in $K_\bullet$. Let Δ denote the comultiplication in $C \otimes_{\hat{\tau}} D$. For any $c \in C^p$ and $d \in D^q$ we have

$$\Delta(c \otimes d) = \sum\nolimits_{r=0}^{p} \sum\nolimits_{u=0}^{q} (-1)^{(p-r)u} c^r_{(1)} \otimes \left(d^u_{(1)}\right)_\tau \otimes \left(c^{p-r}_{(2)}\right)_\tau \otimes d^{q-u}_2,$$

so, the component of $\Delta(c \otimes d)$ in $(C \otimes_{\hat{\tau}} D)^1 \otimes (C \otimes_{\hat{\tau}} D)^{p+q-1}$ is obtained from the above equality by dropping all summands but the ones with either $r = 1$ and $u = 0$, or $r = 0$ and $u = 1$. We get

$$\Delta^{1,p+q-1}(c \otimes d) = \sum c^1_{(1)} \otimes 1 \otimes c^{p-1}_{(2)} \otimes d + (-1)^p \sum\nolimits_{\tau} 1 \otimes \left(d^1_{(1)}\right)_\tau \otimes c_\tau \otimes d^{q-1}_{(2)}.$$

and thus, in degree n, for any $a \in A$, $b \in B$, and any $c \in C^p$, $d \in D^q$ with $p + q = n$ we have:

$$\begin{aligned}\partial_n\left((a \otimes b) \otimes (c \otimes d)\right) &= \sum\nolimits_{\tau} (-1)^p (a \otimes b) \cdot_\sigma \theta(1 \otimes (d^{(1)}_1)_\tau) \otimes c_\tau \otimes d^{(q-1)}_2 + \\ &\quad + \sum (a \otimes b) \cdot_\sigma \theta((c^{(1)}_1) \otimes 1) \otimes c^{(p-1)}_2 \otimes d \\ &= \sum\nolimits_{\tau} (-1)^p a \otimes b\theta_{D,B}((d^{(1)}_1)_\tau) \otimes c_\tau \otimes d^{(q-1)}_2 + \sum a\theta_{C,A}(c^{(1)}_1)_\sigma \otimes b \otimes c^{(p-1)}_2 \otimes d.\end{aligned}$$

To make computations with morphisms in the category of R-bimodules we use string representation of morphisms in a monoidal category, which is explained for example in [Kas95, Chapter XIV.1]. Each morphism will be represented downwards, as a string together with a box. The name of the morphism will be written inside the box. For the identity of a bimodule we shall draw a string without any box. The tensor product and the composition of two morphisms will be represented by horizontal and vertical juxtaposition, respectively. In conclusion, every string diagram may be interpreted as the representation




FIGURE 1. The definition of ∂_n

of a composition $f_1 \circ \cdots \circ f_n$, where each f_i is a tensor product $f_i = \mathrm{Id}_{X_i} \otimes g_i \otimes \mathrm{Id}_{Y_i}$. The corresponding diagrams will be drawn one under the other one, starting with f_1 on the top.

As an example, let us have a look at the picture in Figure 1, which represents ∂_n. The first term is defined on $A^m \otimes B^n \otimes C^p \otimes D^q$ and takes values in $A^m \otimes B^{n+1} \otimes C^p \otimes D^{q-1}$, and can be written (up to a sign) as the composition of four maps

$$(\mathrm{Id}_{A^m} \otimes m_B \otimes \mathrm{Id}_{C^p \otimes D^{q-1}})(\mathrm{Id}_{A^m \otimes B^n} \otimes \theta_{D,B} \otimes \mathrm{Id}_{C^p \otimes D^{q-1}})(\mathrm{Id}_{A^m \otimes B^n} \otimes \tau \otimes \mathrm{Id}_{D^{q-1}})(\mathrm{Id}_{A^m \otimes B^n \otimes C^p} \otimes \Delta^{1,q-1}).$$

As usual, the multiplication of an R-ring is drawn by joining two strings. For the comultiplication of a coring we shall use the 'dual' representation, obtained by turning up-side-down the string diagram of the multiplication of a ring (Δ splits the string representing the domain into two strings). Note that the special notation of σ as a crossing. For τ and λ we shall use the inverse crossing representation, to put stress on the fact they were associated to σ^{-1}, and not σ.

If we define $\delta_n := (\mathrm{Id}_A \otimes \lambda \otimes \mathrm{Id}_D) \circ \partial_n \circ (\mathrm{Id}_A \otimes \lambda^{-1} \otimes \mathrm{Id}_D)$ then we have that $\delta_n = \delta'_n + \delta''_n$, where δ'_n is the first term of ∂_n composed to the left by $\mathrm{Id}_A \otimes \lambda \otimes \mathrm{Id}_D$ and to the right by $\mathrm{Id}_A \otimes \lambda^{-1} \otimes \mathrm{Id}_D$. The map δ''_n is obtained similarly from the second term of ∂_n. The computation of δ'_n is performed in Figure 2. For the first equality we are using (25). The second one means that λ is compatible with the multiplication of B, while the third one is obvious, as λ and λ^{-1} are inverses each other. The morphism δ''_n is computed, using the same method, in Figure 3. To deduce the first identity we are using (26). The compatibility between the coring structure of B and the entwining map λ implies the second equality. The third one is again trivial as λ^{-1} is the inverse of λ. The above computations shows that $\delta'_\bullet = (-1)^\bullet \mathrm{Id}_{\mathrm{K}'_\bullet} \otimes d''$ and $\delta''_\bullet = d' \otimes \mathrm{Id}_{\mathrm{K}''_\bullet}$, where d' and d'' are the differentials of $\mathrm{K}'_\bullet$ and $\mathrm{K}''_\bullet$, respectively. Therefore,$\mathrm{K}^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D) \cong \mathrm{K}^l_\bullet(A, C) \otimes \mathrm{K}^l_\bullet(B, D)$. Let us assume that (A, C) and (B, D) are Koszul. By definition, then the complexes $\mathrm{K}^l_\bullet(A, C)$ and $\mathrm{K}^l_\bullet(B, D)$ are acyclic and their homology groups in degree zero are isomorphic to R. If we prove that $\mathrm{K}^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$ has the same




FIGURE 2. The computation of δ'_n




FIGURE 3. The computation of δ''_n

property, then the augmented complex $\tilde{\mathrm{K}}^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$ would be exact. Since R is semisimple, using Künneth formula [Wei97, Theorem 3.6.3], we get

$$H^n(\mathrm{K}^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)) \cong \bigoplus\nolimits_{p+q=n} H^p(\mathrm{K}^l_\bullet(A, C)) \otimes H^q(\mathrm{K}^l_\bullet(B, D)).$$

Thus $\mathrm{K}^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)$ is clearly acyclic, and

$$H^0(\mathrm{K}^l_\bullet(A \otimes_\sigma B, C \otimes_{\hat{\tau}} D)) \cong H^0(\mathrm{K}^l_\bullet(A, C)) \otimes H^0(\mathrm{K}^l_\bullet(B, D)) \cong R \otimes R \cong R.$$

In conclusion, the theorem is proved. □

Corollary 3.15. *Let $\sigma : B \otimes A \to A \otimes B$ be a twisting map between two Koszul R-rings. Then $A \otimes_\sigma B$ is Koszul and $T(A \otimes_\sigma B) \cong T(A) \otimes_{\hat{\tau}} T(B)$.*

Proof. One applies the preceding theorem, taking τ and λ to be the twisting map and the entwining structures that we constructed in Proposition 3.10. □

We conclude this section by providing an example of two pairs that satisfies the conditions in §3.11. We start with a simple example of Koszul pair.

For every R-bimodule V let $T := T^a_R(V)$. Let $D = R \oplus V$ be the graded coring with comultiplication Δ, such that $\Delta(v) = v \otimes 1 + 1 \otimes v$, for any $v \in V$. We take $\theta_{T,D}$ to be the identity map.

Lemma 3.16. *The pair (T, D) is Koszul.*

Proof. Clearly T and D are graded and connected. The identity from the definition of Koszul pairs is automatically verified, as $D^2 = 0$. Thus (T, D) is pre-Koszul. Furthermore, $\widetilde{\mathrm{K}}^l_\bullet(T, D)$ is the complex

$$0 \longleftarrow R \longleftarrow T \longleftarrow T \otimes V \longleftarrow 0 \longleftarrow \cdots$$

whose non-zero arrows are the projection π^0_T of T on T^0 and d^l_1. Since the multiplication in T is given by juxtaposition of tensor monomials, and $\Delta^{1,0}(v) = v \otimes 1$, for every $v \in V$, we deduce that $d^l_1(x \otimes v) = x \otimes v$. Therefore, d^l_1 identifies $T \otimes V$ and $\overline{T}$. Hence $\widetilde{\mathrm{K}}^l_\bullet(T, D)$ is exact. □

From now on we assume that R is a field and that V has a finite basis $\{e_1, e_2, \ldots, e_n\}$. For a given pre-Koszul pair (A, C), we are going to characterize graded twisting maps $\sigma : T \otimes A \to A \otimes T$, graded twisting maps $\tau : C \otimes D \to D \otimes C$ and graded entwining maps $\lambda : C \otimes T \to T \otimes C$.

3.17. The twisting maps between T and A. Let σ be a twisting map as above. On $T^0 \otimes A$ the map σ is always the canonical identification $T^0 \otimes A \cong A \otimes T^0$. Since σ maps $V \otimes A$ to $A \otimes V$,

$$\sigma(e_i \otimes a) = \sum\nolimits_{j=1}^{n} \sigma_{ij}(a) \otimes e_j,$$

where σ_{ij} is a graded R-linear endomorphisms of A, for every $1 \leq i, j \leq n$. Since σ is compatible with the multiplication of A, it follows that

$$\sigma_{ij}(ab) = \sum\nolimits_{p=1}^{n} \sigma_{ik}(a)\sigma_{kj}(b), \tag{29}$$

for all $a, b \in A$, and all $1 \leq i, j \leq n$. The compatibility with the unit of A is equivalent to the relation

$$\sigma_{ij}(1) = \delta_{i,j}1. \tag{30}$$

Writting down the condition expressing the compatibility between σ and the multiplication of T, we get

$$\sigma(e_{i_1} \cdots e_{i_p} \otimes a) = \sum_{j_1,\dots j_p=1}^{n} \sigma_{i_1 j_1} \cdots \sigma_{i_p j_p}(a) \otimes e_{j_1} \cdots e_{j_p}. \tag{31}$$

In conclusion to every twisting map σ corresponds a matrix $(\sigma_{ij})_{i,j} \in M_n(\mathcal{E}_A)$ such that the relations (29)–(31) hold, where $\mathcal{E}_A$ is the ring of graded linear endomorphisms of A. It is not difficult to see that, conversely, a matrix $(\sigma_{ij})_{i,j}$ satisfying (29) and (30) defines in a unique way a graded twisting map σ. Of course, given $(\sigma_{ij})_{i,j}$, the definition of the corresponding σ is given in (31).

One can prove that σ is invertible if and only if the transposed of the corresponding matrix $(\sigma_{ij})_{i,j}$ is invertible in $M_n(\mathcal{E}_A)$ or, equivalently, $(\sigma_{ij})_{i,j}$ is invertible in $M_n(\mathcal{E}_A^{op})$. More precisely, σ is invertible if and only if there is a matrix $(\sigma'_{ij})_{i,j} \in M_n(\mathcal{E}_A)$ such that $\sum_{j=1}^{n} \sigma_{ji}\sigma'_{kj} = \sum_{j=1}^{n} \sigma'_{ji}\sigma_{kj} = \delta_{i,k}\mathrm{Id}_A$, for all $1 \leq i, k \leq n$. The matrix $(\sigma'_{ij})_{i,j}$ determines the inverse σ' of σ by the formula

$$\sigma'(a \otimes e_{i_1} \cdots e_{i_p}) = \sum_{j_1,\dots j_p=1}^{n} e_{j_1} \cdots e_{j_p} \otimes \sigma'_{i_p j_p} \cdots \sigma'_{i_1 j_1}(a). \tag{32}$$

3.18. The twisting maps between C and D. Now let us fix a graded twisting map $\tau : C \otimes D \to D \otimes C$. Since $D = R \oplus V$ and $\tau(c \otimes 1) = 1 \otimes c$, for all $c \in C$, we have to compute $\tau(c \otimes e_i)$. Let $\mathcal{E}_C$ denote the R-algebra of graded linear endomorphisms of C. Proceeding as in the case of twisting maps of algebras, we see that there is a matrix $(\tau_{ij})_{i,j}$ in $M_n(\mathcal{E}_C)$ such that the following identities hold

$$\tau(c \otimes e_i) = \sum\nolimits_{j=1}^{n} e_j \otimes \tau_{ji}(c), \tag{33}$$

$$\Delta_C\left(\tau_{ij}(c)\right) = \sum\nolimits_{k=1}^{n} \tau_{ik}(c_{(1)}) \otimes \tau_{kj}(c_{(2)}), \tag{34}$$

$$\varepsilon_C\left(\tau_{ij}(c)\right) = \delta_{i,j}\varepsilon_C(c). \tag{35}$$

Conversely, for a matrix that satisfies (34) and (35) , the formula (33) defines a twisting map of coalgebras between C and D. The twisting map τ is invertible if, and only if, the corresponding matrix $(\tau_{ij})_{i,j}$ is invertible in $M_n(\mathcal{E}_C)$. If the inverse is $(\tau'_{ij})_{i,j}$ then τ', the inverse of τ, is given by

$$\tau'(e_i \otimes c) = \sum\nolimits_{j=1}^{n} \tau'_{ji}(c) \otimes e_j. \tag{36}$$

3.19. The entwining map between C and T. Now we are looking for a graded entwining map $\lambda : C \otimes T \to T \otimes C$. The restriction of λ to $C \otimes D$ obviously is a graded twisting map between C and D. Hence, there is a matrix $(\lambda_{ij})_{i,j}$ in $M_n(\mathcal{E}_C)$ such that

$$\lambda(c \otimes e_i) = \sum\nolimits_{j=1}^{n} e_j \otimes \lambda_{ji}(c), \tag{37}$$

$$\Delta_C\left(\lambda_{ij}(c)\right) = \sum\nolimits_{k=1}^{n} \lambda_{ik}(c_{(1)}) \otimes \lambda_{kj}(c_{(2)}), \tag{38}$$

$$\varepsilon_C\left(\lambda_{ij}(c)\right) = \delta_{i,j}\varepsilon_C(c). \tag{39}$$

Furthermore, using the compatibility between λ and the multiplication in T, we get

$$\lambda(c \otimes e_{i_1} \cdots e_{i_p}) = \sum_{j_1,\dots j_p=1}^{n} e_{j_1} \cdots e_{j_p} \otimes \lambda_{j_p i_p} \cdots \lambda_{j_1 i_1}(c). \tag{40}$$

Conversely, if $(\lambda_{ij})_{i,j}$ is a matrix in $M_n(\mathcal{E}_C)$ such that (38) and (39) hold, then the formula (40) defines a graded twisting map λ between C and T, which is invertible if, and only if, the matrix that we start with is so. Note that, if $(\lambda'_{ij})_{i,j}$ is the inverse of $(\lambda_{ij})_{i,j}$, then

$$\lambda'(e_{i_1}\cdots e_{i_p}\otimes c)=\sum_{j_1,\dots j_p=1}^{n}\lambda'_{j_1i_1}\cdots\lambda'_{j_pi_p}(c)\otimes e_{j_1}\cdots e_{j_p}. \tag{41}$$

3.20. The compatibility between σ, τ and λ. We fix a graded twisting map $\sigma : T\otimes A\to A\otimes T$, a graded twisting map $\tau : C\otimes D\to D\otimes C$ and a graded entwining map $\lambda : C\otimes T\to T\otimes C$ as above. Let $(\sigma_{ij})_{i,j}$, $(\tau_{ij})_{i,j}$ and $(\lambda_{ij})_{i,j}$ be the corresponding matrices. We want to explicit the conditions (25) and (26) in this particular setting. For simplicity we assume that $A^1=C^1$ and $\theta_{C,A}=\mathrm{Id}_{C^1}$. Hence the former relation becomes $\lambda^{p,1}=\tau^{p,1}$, for every p. Clearly this is equivalent to the fact that λ and τ are defined by the same matrix in $M_n(\mathcal{E}_C)$. On the other hand, (26) is equivalent to $(\sigma^{q,1})^{-1}=\lambda^{1,q}$, for every q. Taking into the account the fact that the inverse of σ is given by (32), the above identities are equivalent to

$$\sum_{j_1,\dots j_q=1}^{n}e_{j_1}\cdots e_{j_q}\otimes\sigma'_{i_qj_q}\cdots\sigma'_{i_1j_1}(a)=\sum_{j_1,\dots j_q=1}^{n}e_{j_1}\cdots e_{j_q}\otimes\lambda_{j_qi_q}\cdots\lambda_{j_1i_1}(c), \tag{42}$$

for every q and all $c\in C^1$. In the case when $q=1$ we deduce that $\sigma'_{ij}(c)=\lambda_{i,j}(c)$, for all $c\in C^1$ and every $1\le i,j\le n$. Visibly, the fact that $(\sigma'_{ij})_{i,j}$ equals on C^1 the transposed matrix of $(\lambda_{ij})_{i,j}$ implies (42). In conclusion The compatibility relations (25) and (26) are equivalent in our particular setting to the fact that $(\tau_{ij})_{i,j}=(\lambda_{ij})_{i,j}$, and on $C^1=A^1$ we have

$$\sum\nolimits_{j=1}^{n}\sigma_{ji}\lambda_{jk}=\sum\nolimits_{j=1}^{n}\lambda_{ij}\sigma_{kj}=\delta_{i,k}\mathrm{Id}_A, \tag{43}$$

Proposition 3.21. *Let R be a field. Suppose that (A,C) is a pre-Koszul pair such that $A^1=C^1$, and that V is a vector space of dimension n. There is an one-to-one correspondence between the following data:*

(a) The pairs (Σ,Λ), with $\Sigma:=(\sigma_{ij})_{i,j}\in M_n(\mathcal{E}_A)$ and $\Lambda:=(\lambda_{ij})_{i,j}\in M_n(\mathcal{E}_A)$, so that:
 (i) Both Σ^t (the transposed of Σ) and Λ are invertible.
 (ii) The conditions (29) and (30) hold.
 (iii) The conditions (38) and (39) hold.
 (iv) The condition (43) hold on A^1.
(b) The triples (σ,τ,λ) as in §3.11.

The correspondence is given by $(\Sigma,\Lambda)\mapsto(\sigma,\tau,\lambda)$, where σ, τ and λ are defined as in (31), (33) and (40), respectively, where for the definition of τ one takes $\tau_{ij}=\lambda_{ij}$, for all i,j.

Acknowledgments

The third named author was financially supported by CNCSIS, Contract 560/2009 (CNCSIS code ID_69).

References

[BG98] A. Braverman and D. Gaitsgory, *Poincaré-birkhoff-witt theorem for quadratic algebras of koszul type*, J. Algebra **181** (1998), 315–328.

[BGS96] A. Beilinson, V. Ginzburg, and W. Soergel, *Koszul duality patterns in representation theory*, J. Amer. Math. Soc. **9** (1996), 473–527.

[Brz09] T. Brzeziński, *Comodules and corings*, Handbook of Algebra, vol. 6, Elsevier, 2009, pp. 237–318.

[CSV95] A. Cap, H. Schichl, and J. Vanžura, *On twisted tensor products of algebras*, Comm. Algebra **23** (1995), 4701–4735.

[JMLPnPVO] P. Jara Martínez, J. López Peña, F. Panaite, and F. Van Oystaeyen, *On iterated twisted tensor products of algebras*, To appear in *Int. Journal of Math.*

[Kas95] C. Kassel, *Quantum groups*, Graduate Texts in Math., 155, Springer Verlag, Berlin, 1995.

[PP05] A. Polishchuk and L. Positselski, *Quadratic algebras*, University lecture series, vol. 37, American Mathematical Society, Providence, R.I., 2005.

[Pri70] S. Priddy, *Koszul resolutions*, Trans. Amer. Math. Soc. **152** (1970), 39–60.

[VDVK94] A. Van Daele and S. Van Keer, *The Yang–Baxter and Pentagon equation*, Compositio Math. **91** (1994), 201–221.

[Wei97] C. Weibel, *An introduction to homological algebra*, Cambridge University Press, Cambridge, 1997.

University of Granada, Department of Algebra, Granada E-18071, Spain

E-mail address: pjara@ugr.es

Department of Mathematics, University College London, Gower Street, London WC1E 6BT, United Kingdom

E-mail address: jlp@math.ucl.ac.uk

University of Bucharest, Faculty of Mathematics and Computer Science, 14 Academiei Street, Bucharest Ro-010014, Romania

E-mail address: dragos.stefan@fmi.unibuc.ro